2019 Consumer Confidence Report

Water System Name:	Golden Valley Municipal Water District	Report Date: May 2019
		red by state and federal regulations. This report shows the 31, 2019 and may include earlier monitoring data.
Type of water source(s)	in use: Groundwater	
Name & general locatio	on of source(s): Primary Domestic Wel	
Drinking Water Source well with sodium hypo		ter for Gorman community provided by primary domestic
Time and place of regul	arly scheduled board meetings for public pa	articipation: 3 rd Wednesday of each month
For more information, c	contact: Golden Valley Municipal Water	· District Phone: (662) 248-8501

TERMS USED IN THIS REPORT

Maximum Contaminant Level (MCL): The highest level of a contaminant that is allowed in drinking water. Primary MCLs are set as close to the PHGs (or MCLGs) as is economically and technologically feasible. Secondary MCLs are set to protect the odor, taste, and appearance of drinking water.

Maximum Contaminant Level Goal (MCLG): The level of a contaminant in drinking water below which there is no known or expected risk to health. MCLGs are set by the U.S. Environmental Protection Agency (U.S. EPA).

Public Health Goal (PHG): The level of a contaminant in drinking water below which there is no known or expected risk to health. PHGs are set by the California Environmental Protection Agency.

Maximum Residual Disinfectant Level (MRDL): The highest level of a disinfectant allowed in drinking water. There is convincing evidence that addition of a disinfectant is necessary for control of microbial contaminants.

Maximum Residual Disinfectant Level Goal (MRDLG): The level of a drinking water disinfectant below which there is no known or expected risk to health. MRDLGs do not reflect the benefits of the use of disinfectants to control microbial contaminants.

Primary Drinking Water Standards (PDWS): MCLs and MRDLs for contaminants that affect health along with their monitoring and reporting requirements, and water treatment requirements.

Secondary Drinking Water Standards (SDWS): MCLs for contaminants that affect taste, odor, or appearance of the drinking water. Contaminants with SDWSs do not affect the health at the MCL levels.

Treatment Technique (TT): A required process intended to reduce the level of a contaminant in drinking water.

Regulatory Action Level (AL): The concentration of a contaminant which, if exceeded, triggers treatment or other requirements that a water system must follow.

Variances and Exemptions: Permissions from the State Water Resources Control Board (State Board) to exceed an MCL or not comply with a treatment technique under certain conditions.

Level 1 Assessment: A Level 1 assessment is a study of the water system to identify potential problems and determine (if possible) why total coliform bacteria have been found in our water system.

Level 2 Assessment: A Level 2 assessment is a very detailed study of the water system to identify potential problems and determine (if possible) why an *E. coli* MCL violation has occurred and/or why total coliform bacteria have been found in our water system on multiple occasions.

ND: not detectable at testing limit

ppm: parts per million or milligrams per liter (mg/L)

ppb: parts per billion or micrograms per liter (μg/L)

ppt: parts per trillion or nanograms per liter (ng/L)

ppq: parts per quadrillion or picogram per liter (pg/L)

pCi/L: picocuries per liter (a measure of radiation)

The sources of drinking water (both tap water and bottled water) include rivers, lakes, streams, ponds, reservoirs, springs, and wells. As water travels over the surface of the land or through the ground, it dissolves naturally-occurring minerals and, in some cases, radioactive material, and can pick up substances resulting from the presence of animals or from human activity.

Contaminants that may be present in source water include:

- *Microbial contaminants*, such as viruses and bacteria, that may come from sewage treatment plants, septic systems, agricultural livestock operations, and wildlife.
- *Inorganic contaminants*, such as salts and metals, that can be naturally-occurring or result from urban stormwater runoff, industrial or domestic wastewater discharges, oil and gas production, mining, or farming.
- Pesticides and herbicides, that may come from a variety of sources such as agriculture, urban stormwater runoff, and residential uses.
- Organic chemical contaminants, including synthetic and volatile organic chemicals, that are byproducts of
 industrial processes and petroleum production, and can also come from gas stations, urban stormwater runoff,
 agricultural application, and septic systems.
- Radioactive contaminants, that can be naturally-occurring or be the result of oil and gas production and mining activities.

In order to ensure that tap water is safe to drink, the U.S. EPA and the State Board prescribe regulations that limit the amount of certain contaminants in water provided by public water systems. The U.S. Food and Drug Administration regulations and California law also establish limits for contaminants in bottled water that provide the same protection for public health.

Tables 1, 2, 3, 4, 5, and 6 list all of the drinking water contaminants that were detected during the most recent sampling for the constituent. The presence of these contaminants in the water does not necessarily indicate that the water poses a health risk. The State Board allows us to monitor for certain contaminants less than once per year because the concentrations of these contaminants do not change frequently. Some of the data, though representative of the water quality, are more than one year old. Any violation of an AL, MCL, MRDL, or TT is asterisked. Additional information regarding the violation is provided later in this report.

Microbiological Contaminants (complete if bacteria detected)	Highest No. of Detections	No. of Months in Violation	MCL	MCLG	Typical Source of Bacteria
Total Coliform Bacteria (state Total Coliform Rule)	(In a month)		I positive monthly sample ^(a)	0	Naturally present in the environment
Fecal Coliform or <i>E. coli</i> (state Total Coliform Rule)	(In the year)		A routine sample and a repeat sample are total coliform positive, and one of these is also fecal coliform or <i>E. coli</i> positive		Human and animal fecal waste
E. coli (federal Revised Total Coliform Rule)	(In the year)		(b)	0	Human and animal fecal waste

⁽a) Two or more positive monthly samples is a violation of the MCL

⁽b) Routine and repeat samples are total coliform-positive and either is E. coli-positive or system fails to take repeat samples following E. coli-positive routine sample or system fails to analyze total coliform-positive repeat sample for E. coli.

TABLE 2	– SAMPL	ING RESU	LTS SHOW	ING THE D	ETECT	ION O	F LEAD AND C	COPPER
Lead and Copper (complete if lead or copper detected in the last sample set)	Sample Date	No. of Samples Collected	90 th Percentile Level Detected	No. Sites Exceeding AL	AL	РНG	No. of Schools Requesting Lead Sampling	Typical Source of Contaminant
Lead (ppb)	2/26/14 9/29/14 3/2/16 8/23/19	5 5 5 5	<0.002 <0.002 5.1 <0.006	0 0 0 0 0	15	0.2	0	Internal corrosion of household water plumbing systems; discharges from industrial manufacturers; erosion of natural deposits
Copper (ppm)	2/26/14 9/29/14 3/2/16 8/23/19	5 5 5 5	<0.05 <0.05 0.240 0.305	0 0	1.3	0.3	Not applicable	Internal corrosion of household plumbing systems; erosion of natural deposits; leaching from wood preservatives

	TABLE 3	- SAMPLING	RESULTS FOR	SODIUM A	AND HARDI	NESS
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL	PHG (MCLG)	Typical Source of Contaminant
Sodium (ppm)	12/19/07 3/24/14 8/22/17	44 37 37	N/A	None	None	Salt present in the water and is generally naturally occurring
Hardness (ppm)	12/19/07 3/24/14 8/22/17 10/9/19	290 240 220 250	N/A	None	None	Sum of polyvalent cations present in the water, generally magnesium and calcium, and are usually naturally occurring
TABLE 4 – DET	ECTION O	F CONTAMIN	ANTS WITH A	PRIMARY	DRINKING	WATER STANDARD
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
See attached pages 4-12						
TABLE 5 – DETE	CTION OF	CONTAMINA	 NTS WITH A <u>S</u> I	 ECONDAR	<u>Y</u> DRINKIN	G WATER STANDARD
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	SMCL	PHG (MCLG)	Typical Source of Contaminant
See attached page 13						
	TABLE	 6 – DETECTIO	N OF UNREGU	LATED CO	NTAMINA	NTS
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	Notifica	tion Level	Health Effects Language
See attached pages 14-16						

Additional General Information on Drinking Water

Drinking water, including bottled water, may reasonably be expected to contain at least small amounts of some contaminants. The presence of contaminants does not necessarily indicate that the water poses a health risk. More information about contaminants and potential health effects can be obtained by calling the U.S. EPA's Safe Drinking Water Hotline (1-800-426-4791).

Some people may be more vulnerable to contaminants in drinking water than the general population. Immuno-compromised persons such as persons with cancer undergoing chemotherapy, persons who have undergone organ transplants, people with HIV/AIDS or other immune system disorders, some elderly, and infants can be particularly at risk from infections. These people should seek advice about drinking water from their health care providers. U.S. EPA/Centers for Disease Control (CDC) guidelines on appropriate means to lessen the risk of infection by *Cryptosporidium* and other microbial contaminants are available from the Safe Drinking Water Hotline (1-800-426-4791).

Lead-Specific Language: If present, elevated levels of lead can cause serious health problems, especially for pregnant women and young children. Lead in drinking water is primarily from materials and components associated with service lines and home plumbing. Golden Valley Municipal Water District is responsible for providing high quality drinking water, but cannot control the variety of materials used in plumbing components. When your water has been sitting for several hours, you can minimize the potential for lead exposure by flushing your tap for 30 seconds to 2 minutes before using water for drinking or cooking. [OPTIONAL: If you do so, you may wish to collect the flushed water and reuse it for another beneficial purpose, such as watering plants.] If you are concerned about lead in your water, you may wish to have your water tested. Information on lead in drinking water, testing methods, and steps you can take to minimize exposure is available from the Safe Drinking Water Hotline (1-800-426-4791) or at http://www.epa.gov/lead.

TABLE 4 – DETE	ECTION OF CO	NTAMINANT	S WITH A PR	MARY DRI	NKING WAT	TER STANDARD
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
Inorganic Contaminants						
Aluminum (ppm)	3/24/14 8/22/17	0.05 <0.05	0.05 <0.05	1	0.6	Erosion of natural deposits; residue from some surface water treatment processes
Antimony (ppm)	3/24/14 8/22/17	0.002 <0.002	0.002 <0.002	0.006	0.001	Discharge from petroleum refineries; fire retardants; ceramics; electronics; solder
Arsenic (ppb)	12/19/07 8/22/17	2.2 <2.0	2.2 <2.0	10	0.004	Erosion of natural deposits; runoff from orchards; glass and electronics production wastes
Asbestos (MFL)	12/20/14 1/25/16	ND ND	ND ND	7 MFL	7 MFL	Decay of asbestos cement in water mains; erosion of natural deposits
Barium (ppm)	12/19/07 3/24/14 8/22/17	<0.1 0.034 0.034	<0.1 0.034 0.034	1	2	Discharge of oil drilling wastes and from metal refineries; erosion of natural deposits
Beryllium (ppb)	12/19/07 3/24/14 8/22/17	<1.0 1.0 <1.0	<1.0 1.0 <1.0	4	1	Discharge from metal refineries, coal-burning factories, and electrical, aerospace, and defense industries
Cadmium (ppb)	12/19/07 3/24/14 8/22/17	<1.0 1.0 <1.0	<1.0 1.0 <1.0	5	0.04	Internal corrosion of galvanized pipes; erosion of natural deposits; discharge from electroplating and industrial chemical factories and metal refineries; runoff from waste batteries and paints
Chromium (ppb)	12/19/07 3/24/14 8/22/17	<10 10 <10	<10 10 <10	50	(100)	Discharge from steel and pulp mills and chrome plating; erosion of natural deposits
Copper (ppm)	12/19/07 8/22/17 10/9/19	<0.05 <0.01 0.305	<0.05 <0.01 0.305	(1.3)	0.17	Internal corrosion of household plumbing systems erosion of natural deposits; leaching from wood preservatives
Cyanide (ppb)	3/24/14 8/22/17 10/9/19	5 <5 ND	5 <5 ND	150	150	Discharge from steel/metal, plastic and fertilizer factories
Fluoride (ppm)	12/19/07 3/24/14 8/22/17 10/9/19	0.61 0.49 0.75 0.60	0.61 0.49 0.75 0.60	2	1	Erosion of natural deposits; water additive which promotes strong teeth; discharge from fertilizer and aluminum factories
Hexavalent Chromium (ppb)	8/1/18 10/9/19	7.3 1.7	7.3 1.7	10	0.02	Discharge from electroplating factories, leather tanneries, wood preservation, chemical synthesis, refractory production, and textile manufacturing facilities; erosion of natural deposits

TABLE 4 – DETE	ECTION OF CO	NTAMINAN	rs with a pri	MARY DRI	NKING WA'	ΓER STANDARD
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
Lead (ppb)	12/19/07 8/22/17 8/23/19	<2.0 <1.0 <1.0	<2.0 <1.0 <1.0	(AL=15)	0.2	Internal corrosion of household water plumbing systems; discharges from industrial manufacturers; erosion of natural deposits
Mercury (inorganic) (ppb)	12/19/07 3/24/14 8/22/17	<0.20 0.2 <0.20	<0.20 0.2 <0.20	2.0	1.2	Erosion of natural deposits; discharge from refineries and factories; runoff from landfills and cropland
Nickel (ppm)	12/19/07 3/24/14 8/22/17	<0.01 0.01 <0.01	<0.01 0.01 <0.01	0.1	0.012	Erosion of natural deposits; discharge from metal factories
Nitrate (as Nitrogen) (ppm)	6/29/09 3/24/14 3/6/17 8/22/17 8/1/18 12/27/18 10/9/19	4.7 0.05 3.4 3.4 3.2 3.2 3.2	4.7 0.05 3.4 3.4 3.2 3.2 3.2	10	10	Runoff and leaching from fertilizer use; leaching from septic tanks and sewage; erosion of natural deposits
Nitrate (as nitrate, NO ₃) (ppm) 1, 2	1/3/08 4/15/08 6/30/09 9/29/09 3/31/10 7/30/10 3/16/11 12/16/11 3/24/14 4/26/15 6/29/16	35.4 26 21 19 18 20 19 20 16 16	16-35.4	45	45	Runoff and leaching from fertilizer use; leaching from septic tanks and sewage; crosion of natural deposits.
Perchlorate (ppb)	3/24/14 4/26/15 6/30/15 8/25/15 11/18/15 1/25/16 10/9/19	ND ND ND ND ND ND	ND ND ND ND ND ND	6	1	Perchlorate is an inorganic chemical used in solid rocket propellant, fireworks, explosives, flares, matches, and a variety of industries. It usually gets into drinking water as a result of environmental contamination from historic aerospace or other industrial operations that used or use, store, or dispose of perchlorate and its salts.
Selenium (ppb)	12/19/07 3/24/14 8/22/17	2.2 4.0 <2.0	2.2 4.0 <2.0	50	30	Discharge from petroleum, glass, and metal refineries; erosion of natural deposits; discharge from mines and chemical manufacturers; runoff from livestock lots (feed additive)

TABLE 4 – DETI	ECTION OF CO	NTAMINANT	TS WITH A PRI	IMARY DRI	NKING WA	ΓER STANDARD	
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant	
Thallium (ppb)	12/19/07 3/24/14 8/22/17	<1.0 1.0 <1.0	<1.0 1.0 <1.0	2	0.1	Leaching from ore- processing sites; discharge from electronics, glass, and drug factories	
Volatile Organic Contamina	nts			<u> </u>			
Benzene	12/19/07	< 0.50	< 0.50		T	Discharge from plastics, dyes	
(ppb)	4/26/15	ND	ND	1	0.15	and nylon factories; leaching	
A 1-02	6/29/16	ND	ND		9-8000-640-079	from gas storage tanks and	
	8/22/17	< 0.50	< 0.50			landfills	
	8/1/18	ND	ND				
	10/9/19	ND	ND				
Carbon tetrachloride	12/19/07	< 0.50	<0.50	500	100	Discharge from chemical	
(ppt)	4/26/15	ND ND	ND ND			plants and other industrial	
	6/29/16 8/22/17	<0.50	<0.50			activities	
	8/1/18	ND	ND				
	10/9/19	ND	ND				
1, 2-Dichlorobenzene	12/19/07	< 0.50	< 0.50	600	600	Discharge from industrial	
(ppb)	4/26/15	ND	ND			chemical factories	
67 4. 4 6	6/29/16	ND	ND				
	8/22/17	< 0.50	< 0.50				
	8/1/18	ND	ND				
1.18:11	10/9/19	ND 10.50	ND 10.50	ļ <u>-</u>			
1,4-Dichlorobenzene	12/19/07 4/26/15	<0.50 ND	<0.50 ND	5	6	Discharge from industrial chemical factories	
(ppb)	6/29/16	ND ND	ND ND			chemical factories	
	8/22/17	< 0.50	<0.50				
	8/1/18	ND	ND				
	10/9/19	ND	ND				
1, 1-Dichloroethane	12/19/07	< 0.50	< 0.50	5	N/A	Extraction and degreasing	
(ppb)	4/26/15	ND	ND			solvent; used in the	
	6/29/16	ND	ND			manufacture of	
	8/22/17	< 0.50	< 0.50			pharmaceuticals, stone, clay,	
	8/1/18	ND	ND			and glass products; fumigant	
1, 2-Dichloroethane	10/9/19 12/19/07	ND <0.50	ND	500	400	Discharge from in test 1	
(ppt)	4/26/15	<0.50 ND	<0.50 ND	300	400	Discharge from industrial chemical factories	
(PPC)	6/29/16	ND	ND ND			chemical factories	
	8/22/17	< 0.50	< 0.50				
	8/1/18	ND	ND				
	10/9/19	ND	ND				
1, 1-Dichloroethene	12/19/07	< 0.50	< 0.50	6	10	Discharge from industrial	
(ppb)	4/26/15	ND	ND			chemical factories	
	6/29/16	ND	ND				
	8/22/17	<0.50	<0.50				
	8/1/18 10/9/19	ND ND	ND ND				
cis-1,2-Dichloroethene	12/19/07	<0.50	<0.50	6	70	Discharge from industrial	
(ppb)	4/26/15	<0.50 ND	<0.50 ND	0	/0	Discharge from industrial chemical factories; major	
(Ppo)	6/29/16	ND	ND ND			biodegradation byproduct	
	8/22/17	< 0.50	< 0.50		1	of TCE and PCE	
	8/1/18	ND	ND		1	groundwater contamination	
	10/9/19	ND	ND				

TABLE 4 – DETI	ECTION OF CO	NTAMINAN	ΓS WITH A PRI	IMARY DRI	NKING WA	TER STANDARD
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
trans-1,2-Dichloroethene (ppb)	12/19/07 4/26/15 6/29/16 8/22/17 8/1/18 10/9/19	<0.50 ND ND <0.50 ND ND	<0.50 ND ND <0.50 ND ND	10	100	Discharge from industrial chemical factories; minor biodegradation byproduct of TCE and PCE groundwater contamination
Methylene Chloride (Dichloromethane) (ppb)	6/29/16 8/22/17 8/1/18 10/9/19	ND <0.50 ND ND	ND <0.50 ND ND	5	4	Discharge from pharmaceutical and chemica factories; insecticide
1,2-Dichloropropane (ppb)	12/19/07 4/26/15 6/29/16 8/22/17 8/1/18 10/9/19	<0.50 ND ND <0.50 ND	ND <0.50 ND <0.50 ND ND ND	5	0.5	Discharge from industrial chemical factories; primary component of some fumigants
1,3-Dichloropropene (ppt)	12/19/07 4/26/15 6/29/16 8/22/17 8/1/18 10/9/19	<0.50 ND ND <0.50 ND ND	<0.50 ND ND <0.50 ND ND	500	200	Runoff/leaching from nematocide used on cropland
Ethylbenzene (ppb)	12/19/07 4/26/15 6/29/16 8/22/17 8/1/18 10/9/19	<0.50 ND ND <0.50 ND ND	<0.50 ND ND <0.50 ND ND	300	300	Discharge from petroleum refineries; industrial chemica factories
Methyl-tert-butyl ether (ppb)	4/26/15 6/29/16 8/22/17 8/1/18 10/9/19	ND ND <0.50 ND ND	ND ND ND <0.50 ND ND	13	13	Leaking underground storage tanks; discharges from petroleum and chemical factories
Chlorobenzene (ppb)	6/29/16 8/22/17 8/1/18 10/9/19	ND <0.50 ND ND	ND <0.50 ND ND	70	70	Discharge from industrial an agricultural chemical factoric and drycleaning facilities
Styrene (ppb)	4/26/15 6/29/16 8/22/17 8/1/18 10/9/19	ND ND <0.50 ND ND	ND ND <0.50 ND ND	100	0.5	Discharge from rubber and plastic factories; leaching from landfills
1,1,2,2-Tetrachloroethane (ppb)	12/19/07 4/26/15 6/29/16 8/22/17 8/1/18 10/9/19	ND <0.50 ND <0.50 ND ND	ND <0.50 ND <0.50 ND ND	I	0.1	Discharge from industrial and agricultural chemical factories; solvent used in production of TCE, pesticides, varnish and lacquers
Tetrachloroethene (PCE) (ppb)	12/19/07 6/29/16 8/22/17 8/1/18 10/9/19	<0.50 ND <0.50 ND ND	<0.50 ND <0.50 ND ND	5	0.06	Discharge from factories, dry cleaners, and auto shops (metal degreaser)

Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
1,2,4-Trichlorobenzene	12/19/07	< 0.50	<0.50	5	5	Discharge from textile-
(ppb)	6/29/16	ND	ND			finishing factories
	8/22/17	< 0.50	< 0.50			
	8/1/18	ND	ND			
	10/9/19	ND	ND			
1,1,1-Trichloroethane	12/19/07	<0.50	< 0.50	200	1000	Discharge from metal
(ppb)	6/29/16	ND	ND	***		degreasing sites and other
	8/22/17 8/1/18	<0.50 ND	<0.50 ND			factories; manufacture of foo
	10/9/19	ND ND	ND ND	***************************************		wrappings
1,1,2 – Trichloroethane	12/19/07	<0.50	<0.50	5	0.3	Discharge from industrial
(ppb)	6/29/16	ND	ND ND		0.5	chemical factories
(рро)	8/22/17	< 0.50	< 0.50	-		chemical factories
	8/1/18	ND	ND ND			
	10/9/19	ND	ND			
Trichlorofluoromethane	12/19/07	< 0.50	< 0.50	5	1.7	Discharge from metal
(ppb)	6/29/16	ND	ND		''	degreasing sites and other
	8/22/17	< 0.50	< 0.50			factories
	8/1/18	ND	ND			
	10/9/19	ND	ND			
1,1,2-Trichloro-1,2,2	12/19/07	< 0.50	< 0.50	1.2	4	Discharge from metal
trifluoroethane	6/29/16	ND	ND			degreasing sites and other
(ppm)	8/22/17	< 0.50	< 0.50			factories; drycleaning solven
	8/1/18	ND	ND			refrigerant
	10/9/19	ND	ND			
Trichloroethene (TCE)	12/19/07	< 0.50	< 0.50	5	1.7	Discharge from metal degreasing sites and other factories
(ppb)	6/29/16	ND	ND			
	8/22/17	< 0.50	<0.50			
	8/1/18	ND ND	ND ND			
Toluene	10/9/19 4/26/15	ND ND	ND ND	150	150	Discharge from petroleum
(ppb)	12/19/07	< 0.50	<0.50	130	130	and chemical factories;
(ppo)	6/29/16	ND	ND			underground gas tank
	8/22/17	< 0.50	<0.50			leaks
	8/1/18	ND	ND			lettes
	10/9/19	ND	ND			
Vinyl Chloride	12/19/07	ND	ND	500	50	Leaching from PVC piping;
(ppt)	4/26/15	< 0.50	< 0.50			discharge from plastics
	6/29/16	ND	ND			factories; biodegradation
	8/22/17	< 0.50	< 0.50			byproduct of TCE and PCE
	8/1/18	ND	ND			groundwater contamination
	10/9/19	ND	ND			
Total Xylenes	12/19/07	ND	ND	1.750	1.8	Discharge from petroleum
(ppm)	4/26/15	< 0.50	< 0.50		4	and chemical factories; fuel
	6/29/16	ND	ND			solvent
	8/22/17	< 0.50	<0.50			
	8/1/18	ND	ND			
0 371	10/9/19	ND	ND	 	<u> </u>	T21 1
- & m-Xylenes	6/29/16	ND	ND			Discharge from petroleum an
	8/22/17 8/1/18	<0.50 ND	<0.50	1	1	chemical factories; fuel solvent
	10/9/19	ND ND	ND ND			
-Xylene	6/29/16	ND ND	ND ND			Discharge from petroleum an
-Aylene	8/22/17	<0.50	<0.50	1		chemical factories; fuel
	8/1/18	ND	ND			solvent
	10/9/19	ND ND	ND ND			Borront
	10/2/12	1127	1,12	1		
			·			
	1		I	1		I

TABLE 4 – DETE	CTION OF CO	NTAMINANT	S WITH A PR	IMARY DRI	NKING WA	TER STANDARD
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
Synthetic Organic Contamina	ents including Pes	ticides and Her	bicides			
2,4-D (ppb)	4/26/15 5/18/18	ND ND	ND ND	70	20	Runoff from herbicide used or row crops, range land, lawns, and aquatic weeds
2,4,5-TP (Silvex) (ppb)	4/26/15 5/18/18	ND ND	ND ND	50	3	Residue of banned herbicide
Alachlor (ppb)	4/26/15 5/18/18	ND ND	ND ND	2	4	Runoff from herbicide used or row crops
Atrazine (ppb)	4/26/15 5/18/18	ND ND	ND ND	1	0.15	Runoff from herbicide used on row crops and along railroad and highway right- of-ways
Bentazon (ppb)	4/26/15 5/18/18	ND ND	ND ND	18	200	Runoff/leaching from herbicide used on beans, peppers, corn, peanuts, rice, and ornamental grasses
Benzo(a)pyrene (PAH) (ppt)	4/26/15 5/18/18	ND ND	ND ND	200	7	Leaching from linings of wate storage tanks and distribution mains
Carbofuran (ppb)	4/26/15 5/18/18	ND ND	ND ND	18	1.7	Leaching of soil fumigant use on rice and alfalfa, and grape vineyards
Chlordane (ppt)	5/18/18	ND	ND	100	30	Residue of banned insecticide
Dalapon (ppb)	4/26/15 5/18/18	ND ND	ND ND	200	790	Runoff from herbicide used or rights-of-ways, and crops and landscape maintenance
Di(2-ethylhexyl) adipate (ppb)	4/26/15 5/18/18	ND ND	ND ND	400	200	Discharge from chemical factories
Di(2-ethylhexyl) phthalate (ppb)	5/18/18	ND	ND	4	12	Discharge from rubber and chemical factories; inert ingredient in pesticides
Dibromochloropropane (DBCP) (pptb)	4/26/15 5/18/18	ND ND	ND ND	200	1.7	Banned nematocide that may still be present in soils due to runoff/leaching from former use on soybeans, cotton, vineyards, tomatoes, and tree fruit
Dinoseb (ppb)	4/26/15 5/18/18	ND ND	ND ND	7	14	Runoff from herbicide used or soybeans, vegetables, and fruits
Dioxin (2,3,7,8-TCDD) (ppq)	4/26/15 5/18/18	ND ND	ND ND	30	0.05	Emissions from waste incineration and other combustion; discharge from chemical factories
Diquat (ppb)	4/26/15 5/18/18	ND ND	ND ND	20	15	Runoff from herbicide use for terrestrial and aquatic weeds

SWS CCR Form

DUC									
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant			
Endothal (ppb)	4/26/15 5/18/18	ND ND	ND ND	100	94	Runoff from herbicide use for terrestrial and aquatic weeds; defoliant			
Endrin (ppb)	5/18/18	ND	ND	2	1.8	Residue of banned insecticide and rodenticide			
Ethylene dibromide (EDB) (ug/L)	4/26/15 9/9/15 5/18/18	ND ND ND	ND ND	50	10	Discharge from petroleum refineries; underground gas tank leaks; banned nematocid that may still be present in soils due to runoff and leaching from grain and fruit crops			
Glyphosate (ppb)	4/26/15 5/18/18	ND ND	ND ND	700	900	Runoff from herbicide use			
Heptachlor (ppt)	4/26/15 5/18/18	ND ND	ND ND	10	8	Residue of banned insecticide			
Heptachlor epoxide (ppt)	4/26/15 5/18/18	ND ND	ND ND	10	6	Breakdown of heptachlor			
Hexachlorobenzene (ppb)	4/6/15 5/18/18	ND ND	ND ND	1	0.03	Discharge from metal refineries and agricultural chemical factories; byproduct of chlorination reactions in wastewater			
Hexachlorocyclopentadiene (ppb)	4/26/15 5/18/18	ND ND	ND ND	50	2	Discharge from chemical factories			
Lindane (ppt)	4/26/15 5/18/18	ND ND	ND ND	200	32	Runoff/leaching from insecticide used on cattle, lumber, and gardens			
Methoxychlor (ppb)	4/26/15	ND	ND	30	0.09	Runoff/leaching from insecticide used on fruits, vegetables, alfalfa, and livestock			
Molinate (Ordram) (ppb)	4/26/15 5/18/18	ND ND	ND ND	20	1	Runoff/leaching from herbicide used on rice			
Oxamyl (Vydate) (ppb)	4/26/15 5/18/18	ND ND	ND ND	50	26	Runoff/leaching from insecticide used on field crops fruits and ornamentals, especially apples, potatoes, and tomatoes			
PCBs (Polychlorinated biphenyls) (ppt)	4/26/15 5/18/18	ND ND	ND ND	500	90	Runoff from landfills; discharge of waste chemicals			
Pentachlorophenol (ppb)	4/26/15 5/18/18	ND ND	ND ND	1	0.3	Discharge from wood preserving factories, cotton and other insecticidal/herbicidal uses			
Picloram (ppb)	4/26/15 5/18/18	ND ND	ND ND	500	500	Herbicide runoff			

Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
Simazine (ppb)	4/26/15 5/18/18	ND ND	ND ND	4	4	Herbicide runoff
Thiobencarb (ppb)	4/26/15 5/18/18	ND ND	ND ND	70	70	Runoff/leaching from herbicide used on rice
Toxaphene (ppb)	4/26/15 5/18/18	ND ND	ND ND	3	0.03	Runoff/leaching from insecticide used on cotton and cattle
Radioactive Contaminants						
Gross Beta Particle Activity (pCi/L)	6/30/15	4.60	4.60	50 (a)	(0)	Decay of natural and man- made deposits
(a) Effective 6/11/2006, the gross beta particle activity MCL is 4 millirems/year annual dose equivalent to the total body or any internal organ. 50 pCi/L is used as a screening level.	=		÷.			=
Strontium-90 (pCi/L)				8	0.35	Decay of natural and man- made deposit
Tritium (pCi/L)				20,000	400	Decay of natural and man- made deposits
Gross Alpha Particle Activity (pCi/L)	4/26/15 6/30/15 8/25/15	4.60 4.60 1.28	±0.348 ±0.465	15	(0)	Erosion of natural deposits
Radium 226 (pCi/L)	4/26/15 6/30/15 8/25/15 11/17/15 1/25/16	0.278±0.546 (0.981) 0.0993±0.435 (0.922) 0.0798±0.271 (0.588) 1.97±1.02 (0.944) 0.528±0.606 (0.946)	0.278±0.546 (0.981) 0.0993±0.435 (0.922) 0.0798±0.271 (0.588) 1.97±1.02 (0.944) 0.528±0.606 (0.946)	5	(O)(b)	Erosion of natural deposits
Radium 228 (pCi/L)	4/26/15 6/30/15	0.517±0.384 (0.773) 0.385±0.425 (0.902) 0.0574±0.363 (0.710) 0.839±0.44 (0.842) 0.399±0.365 (0.758)	0.517±0.384 (0.773) 0.385±0.425 (0.902) 0.0574±0.363 (0.710) 0.839±0.44 (0.842) 0.399±0.365 (0.758)	5	(0)(6)	Erosion of natural deposits
Combined Radium-226 and Radium-228 (pCi/L)	6/29/16	1.34±1.08 (1.80)	1.34±1.08 (1.80)	5	n/a	Erosion of natural deposits

Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	MCL [MRDL]	PHG (MCLG) [MRDLG]	Typical Source of Contaminant
(b) If reporting results for Ra-226 and Ra-228 as individual constituents, the PHG is 0.05 pCi/L for Ra-226 and 0.019 pCi/L for Ra-228.						
Uranium (pCi/L)	3/24/14 4/26/15	5.4 4.9	5.4 4.9	20	0.43	Erosion of natural deposits
Disinfection Byproducts, Disin	nfectant Residuals	s, and Disinfect	ion Byproduct Pr	ecursors		1-1-1-1
TTHMs (Total Trihalomethanes) (ppb)	11/11/08 3/23/09 9/29/09 3/31/10 7/30/10 3/16/11 12/16/11 8/1/18 10/9/19	12.0 13.0 8.8 9.0 4.2 8.7 8.3 14.0 3.8	4.2 – 14.0	80	N/A	By-product of drinking water disinfection
Halogenated Acetic Acids (HAA5) (ppb)	11/11/08 3/23/09 9/29/09 3/31/10 7/30/10 3/16/11 12/16/11 8/22/17 8/1/18 10/9/19	2.5 3.7 2.8 3.6 2.5 3.4 2.0 1.7 1.4 ND	ND – 3.7	60	N/A	Byproduct of drinking water disinfection
Bromate (ppb)				10	0.1	Byproduct of drinking water disinfection
Chloramines (ppm)				[MRDL = 4.0 (as Cl ₂)]	[MRDLG = 4 (as Cl ₂)]	Drinking water disinfectant added for treatment
Chlorine (ppm)		ts		[MRDL = 4.0 (as Cl ₂₎]	[MRDLG = 4 (as Cl ₂₎	Drinking water disinfectant added for treatment
Chlorite (ppm)				1.0	0.05	Byproduct of drinking water disinfection
Chlorine Dioxide (ppb)				[MRDL = 800 (as ClO ₂)]	[MRDLG = 800 (as ClO ₂)]	Drinking water disinfectant added for treatment
Control of DBP precursors (TOC)				TT	N/A	Various natural and man-mad sources

Chemical or Constituent	Unit Measurement	Sample Date	Level Detected	MCL	Typical Source of Contaminant
Aluminum	ррЬ	12/19/07	<50	200	Erosion of natural deposits; residual from some surface water treatment processes
Color	Units	12/19/07 3/24/14 8/22/17 10/9/19	<3.0 1.0 1.0 2.0	15	Naturally-occurring organic materials
Copper	ppm	12/19/07 3/24/14 8/22/17	1.0 0.01 <0.01	1,0	Internal corrosion of household plumbing systems; erosion of natural deposits; leaching from wood preservatives
Foaming Agents (MBAS)	ppm	12/19/07 3/24/14 8/22/17 10/9/19	<0.050 0.10 <0.10 ND	500	Municipal and industrial waste discharges
Iron	ppb	12/19/07 3/24/14 8/22/17	<100 50 <50	300	Leaching from natural deposits; industrial wastes
Manganese	ppb	12/19/07 3/24/14 3/6/17 8/22/17	<20 10 ND <10	50	Leaching from natural deposits
Methyl-tert-butyl ether (MTBE)	ppb	12/19/07 3/24/14	NT ND	5	Leaking underground storage tanks; discharge from petroleum and chemical factories
OdorThreshold	Units	12/19/07 3/24/14 8/22/17 10/9/19	<1.0 0 0 0	3	Naturally-occurring organic materials
Silver	ppb	12/19/07 3/24/14 8/22/17	<10 10 <10	100	Industrial discharges
Thiobencarb	ppb		NT	1	Runoff/leaching from rice herbicide
Turbidity	NFU	12/19/07 3/24/14 8/22/17 10/9/19	0.16 0.1 0.28 ND	5	Soil runoff
Zinc	ppm	12/19/07 3/24/14 8/22/17	<0.05 0.05 <0.05	5.0	Runoff/leaching from natural deposits; industrial wastes
Total Dissolved Solids (TDS)	ppm	12/19/07 3/24/14 8/22/17 10/9/19	430 430 400 420	1000	Runoff/leaching from natural deposits
Specific Conductance/Electrical Conductivity	μmhos/cm	12/19/07 3/17/11 3/24/14 8/22/17 10/9/19	600 600 572 593 622	1600	Substances that form ions when in water; seawater influence

TABLE 5 – DETEC	TION OF CONTAI	MINANTS WITH	A SECONDARY DR	INKING WA	TER STANDARD
Chemical or Constituent	Unit Measurement	Sample Date	Level Detected	MCL	Typical Source of Contaminant
Chloride	ppm	12/19/07 3/24/14 8/22/17 10/9/19	21 15 15 24	500	Runoff/leaching from natural deposits; seawater influence
Sulfate .	ppm	12/19/07 3/24/14 8/22/17 10/9/19	71 70 71 75	500	Runoff/leaching from natural deposits; industrial wastes

	TABLE	6 – DETECTION	OF UNREGUI	LATED CONTAMINA	ANTS
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	Notification Level	Health Effects Language
1,3-Dichloropropene (ug/L)	12/19/07 8/22/17	<0.50 <0.50	<0.50 <0.50	N/A	Some people who use water containing 1,3-dichloropropene in excess of the MCL over many years may have an increased risk of getting cancer.
1,3-Dichlorobenzene (ug/L)	12/19/07	<0.50	<0.50	N/A	Causes eye and skin irritation. May be absorbed through the skin. Causes gastrointestinal irritation with nausea, vomiting and diarrhea. Chronic exposure may cause liver and kidney damage.
1,3,5-Trimethylbenzene (TMB) (ug/L)	12/19/07	<0.50	<0.50	N/A	Short-term health effects include irritation and burning of the skin and eyes. Long-term health effects include liver damage, anemia, and respiratory effects.
1,3-Dichloropropane (ug/L)	12/19/07	<0.50	<0.50	N/A	Acute inhalation exposure effects include mucous membrane irritation, chest pain, and breathing difficulties. Chronic dermal exposure may result in skin sensitization. May cause damage to the nasal mucosa and urinary bladder. Classified as a probable human carcinogen.
1,2,4-Trimethlybenzene (ug/L)	12/19/07	<0.50	<0.50	N/A	Inhalation causes confusion, cough, dizziness, drowsiness, headache, sore throat, and vomiting. Causes dry, red, itchy skin, and painful red eyes.
Sec-Butylbenzene (ug/L)	12/19/07	<0.50	<0.50	N/A	Causes eye irritation. Causes skin irritation. May cause gastrointestinal irritation with nausea, vomiting and diarrhea. Ingestion of large amounts may cause central nervous system (CNS) depression. Causes respiratory tract irritation. Prolonged or repeated skin contac may cause dermatitis.

Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	Notification Level	Health Effects Language
1,2,3-Trichlorobenzene (ug/L)	12/19/07	<0.50	<0.50	N/A	Potential acute effects include changes in liver, kidneys and adrenal glands. Long-term (chronic) exposure potentially causes increased adrenal gland weights.
2-Chloroethlylvinylether (ug/L)	12/19/07	<0.50	<0.50	N/A	Dangerous when exposed to heat, flame, or oxidizers. Produces eye irritation, skin irritation. When heated, may emit highly toxic vapors.
Tributyl Alcohol, TBA (ug/L)	12/19/07	<10	<10	N/A	Produces eye irritation, skin irritation. If inhaled, can cause nose, throat, and lung irritation. Can cause headaches, dizziness, lightheadedness, and unconsciousness.
1,1,1,2-Tetrachloroethane (ug/L)	12/19/07	<0.50	<0.50	N/A	Can cause shallow breathing, faint pulse, decreased blood pressure, and possibly unconsciousness. Possible carcinogen.
1,2,3-Trichloropropane (ug/L)	12/19/07 9/20/18 11/29/18	<0.50 ND ND	<0.50 ND ND	N/A	Classified as a carcinogen. Enters your body when drinking water or breathing air containing this chemical. The human body absorbs much or all of it that is in drinking water. Exposure can also occur by inhalation within the household (such as in the shower).
Bromomethane (ug/L)	12/19/07	<0.50	<0.50	N/A	Inhalation causes headaches, weakness and nausea. Inhaling a large amount causes fluid to build up in the lungs. May result in muscle tremors, seizures, decreased kidney function, and urine production may slow or stop. Swallowing it causes stomach irritation. Skin exposure causes itchy, red skin and blisters.
Ethyl-tert-Butyl Ether, ETBE (ug/L)	12/19/07	<0.50	<0.50	N/A	N/A
Dichlorodifluoromethane (ug/L)	12/19/07	<0.50	<0.50	N/A	Consuming large amounts of dichlorodifluoromethane may cause neurological and cardiac effects. Long- term exposures to dichlorodifluoromethane resulted in smaller body weight in laboratory animals.
Dibromomethane (ug/L)	12/19/07	<0.50	<0.50	N/A	Can affect the brain, damage skin, damage sperm in males, and even cause death if exposure is very high.
Isopropylbenzene (ug/L)	12/19/07	<0.50	<0.50	N/A	Causes dizziness, ataxia, drowsiness, headache, and potentially unconsciousness.

	TABLE 6 – DETECTION OF UNREGULATED CONTAMINANTS							
Chemical or Constituent (and reporting units)	Sample Date	Level Detected	Range of Detections	Notification Level	Health Effects Language			
Di-Isopropyl Ether, DIPE (ug/L)	12/19/07	<0.50	<0.50	N/A	N/A			
Chloromethane (ug/L)	12/19/07	<0.50	<0.50	N/A	Exposure to large amounts may cause liver problems and may result in an increased risk of getting cancer.			
Chloroethane (ug/L)	12/19/07	<0.50	<0.50	N/A	Exposure to high concentrations can affect your nervous system, causing lack of muscle control and unconsciousness.			
n-Butylbenzene (ug/L)	12/19/07	<0.50	<0.50	N/A	N/A			

^{*}Any violation of an MCL, MRDL, or TT is asterisked. Additional information regarding the violation is provided later in this report

Additional General Information on Drinking Water

Drinking water, including bottled water, may reasonably be expected to contain at least small amounts of some contaminants. The presence of contaminants does not necessarily indicate that the water poses a health risk. More information about contaminants and potential health effects can be obtained by calling the U.S. EPA's Safe Drinking Water Hotline (1-800-426-4791).

Some people may be more vulnerable to contaminants in drinking water than the general population. Immunocompromised persons such as persons with cancer undergoing chemotherapy, persons who have undergone organ transplants, people with HIV/AIDS or other immune system disorders, some elderly, and infants can be particularly at risk from infections. These people should seek advice about drinking water from their health care providers. U.S. EPA/Centers for Disease Control (CDC) guidelines on appropriate means to lessen the risk of infection by *Cryptosporidium* and other microbial contaminants are available from the Safe Drinking Water Hotline (1-800-426-4791).

Lead-Specific Language for Community Water Systems: If present, elevated levels of lead can cause serious health problems, especially for pregnant women and young children. Lead in drinking water is primarily from materials and components associated with service lines and home plumbing. [INSERT NAME OF UTILITY] is responsible for providing high quality drinking water, but cannot control the variety of materials used in plumbing components. When your water has been sitting for several hours, you can minimize the potential for lead exposure by flushing your tap for 30 seconds to 2 minutes before using water for drinking or cooking. [Optional: If you do so, you may wish to collect the flushed water and reuse it for another beneficial purpose, such as watering plants.] If you are concerned about lead in your water, you may wish to have your water tested. Information on lead in drinking water, testing methods, and steps you can take to minimize exposure is available from the Safe Drinking Water Hotline (1-800-426-4701) or at http://www.epa.gov/lead.

Pa	ge	1	7	0	f	1	9	

Summary Information for Violation of a MCL, MRDL, AL, TT, or Monitoring and Reporting Requirement

VIOLATION OF A MCL, MRDL, AL, TT, OR MONITORING AND REPORTING REQUIREMENT							
Violation	Explanation	Duration	Actions Taken to Correct the Violation	Health Effects Language			

For Water Systems Providing Groundwater as a Source of Drinking Water

TABLE 7 – SAMPLING RESULTS SHOWING FECAL INDICATOR-POSITIVE GROUNDWATER SOURCE SAMPLES								
Microbiological Contaminants (complete if fecal-indicator detected) Total No. of Detections Sample Dates MCL [MRDL] [MRDL] Typical Source of Contaminant								
E. coli	(In the year)		0	(0)	Human and animal fecal waste			
Enterococci	(In the year)		TT	N/A	Human and animal fecal waste			
Coliphage	(In the year)		ТТ	N/A	Human and animal fecal waste			

Summary Information for Fecal Indicator-Positive Groundwater Source Samples, Uncorrected Significant Deficiencies, or Groundwater TT

SPECIAL	NOTICE OF FECAL IN	DICATOR-POSITIVE	GROUNDWATER SOURCE S	AMPLE
	SPECIAL NOTICE FOR	UNCORRECTED SIG	NIFICANT DEFICIENCIES	
	VIOL	ATION OF GROUNDW	ATER TT	
TT Violation	Explanation	Duration	Actions Taken to Correct the Violation	Health Effects Language

For Sy	stems Providing Si	ırface Water as a	Source of Drinking Wa	iter
TABLE 8 - SA	AMPLING RESULTS SI	HOWING TREATME	ENT OF SURFACE WATER SO	OURCES
atment Technique (a) see of approved filtration tec	chnology used)			keparti kelekera da minga kelekera kelekera para sara a na mana da mana sara da mana sara da mana sara da mana
pidity Performance Standar t must be met through the v	·ds ^(b)	2 – Not exceed	lered water must: equal to NTU in 95% of meas NTU for more than eight consecu _ NTU at any time.	
vest monthly percentage of formance Standard No. 1.	samples that met Turbidity	San Assessment		
hest single turbidity measur	rement during the year			
Turbidity (measured in NT Turbidity results which me	d to reduce the level of a con U) is a measurement of the et performance standards are	cloudiness of water and considered to be in component on for Violation (is a good indicator of water quality a liance with filtration requirements. of a Surface Water TT	and filtration perform
irements A required process intended Furbidity (measured in NT Furbidity results which me	d to reduce the level of a con U) is a measurement of the et performance standards are	cloudiness of water and considered to be in comp	is a good indicator of water quality a liance with filtration requirements. of a Surface Water TT	Health Effects
irements A required process intended Furbidity (measured in NT Furbidity results which me	d to reduce the level of a con (U) is a measurement of the et performance standards are Immary Information VIOLATI	cloudiness of water and considered to be in component on for Violation (ION OF A SURFACE	is a good indicator of water quality a diance with filtration requirements. of a Surface Water TT WATER TT Actions Taken to Correct	

Consumer Confidence Report	Page 19 of 19
Summary Information for Federal Revised Total Coliform Ru	ile
Level 1 and Level 2 Assessment Requirements	
Level 1 or Level 2 Assessment Requirement not Due to an <i>E. coli</i> MCL V	liolation
Coliforms are bacteria that are naturally present in the environment and are used as an indicato harmful, waterborne pathogens may be present or that a potential pathway exists through which could the drinking water distribution system. We found coliforms indicating the need to look for pote treatment or distribution. When this occurs, we are required to conduct assessment(s) to identify any problems that were found during these assessments.	ontamination may enter ntial problems in water
During the past year we were required to conduct 0 Level 1 assessment(s). Zero Level 1 assessmen addition, we were required to take 0 corrective actions and we completed 0 of these actions.	t(s) were completed. In
During the past year 0 Level 2 assessments were required to be completed for our water system. Zo were completed. In addition, we were required to take 0 corrective actions and we completed 0 of	
Level 2 Assessment Requirement Due to an E. coli MCL Violation	1
E. coli are bacteria whose presence indicates that the water may be contaminated with human or pathogens in these wastes can cause short-term effects, such as diarrhea, cramps, nausea, headach They may pose a greater health risk for infants, young children, the elderly, and people with severely systems. We found E. coli bacteria, indicating the need to look for potential problems in water tr When this occurs, we are required to conduct assessment(s) identify problems and to correct any produring these assessments.	nes, or other symptoms. y-compromised immune reatment or distribution.
We were required to complete a Level 2 assessment because we found <i>E. coli</i> in our water system required to take 0 corrective actions and we completed 0 of these actions.	n. In addition, we were

APPENDIX A: Regulated Contaminants with Primary Drinking Water Standards

Microbiological Contaminants

Contaminant	Unit Measure	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Total Coliform Bacteria (state Total Coliform Rule)	MCL: Systems that collect more samples per m 5.0% of monthly sam are positive Systems that collect than 40 samples per month:	MCL: Systems that collect 40 or more samples per month: 5.0% of monthly samples are positive Systems that collect less than 40 samples per month:	(0)	Naturally present in the environment	Coliforms are bacteria that are naturally present in the environment and are used as an indicator that other, potentially-harmful, bacteria may be present. Coliforms were found in more samples than allowed and this was a warning of potential problems.
Fecal coliform and <i>E. coli</i> (state Total Coliform Rule)	MCL: A routine sample and a repeat sample are total coliform positive, and of these is also fecal coliffor or E. coli positive	MCL: A routine sample and a repeat sample are total coliform positive, and one of these is also fecal coliform or <i>E. coli</i> positive	(0)	Human and animal fecal waste	Fecal coliforms and <i>E. coli</i> are bacteria whose presence indicates that the water may be contaminated with human or animal wastes. Microbes in these wastes can cause short-term effects, such as diarrhea, cramps, nausea, headaches, or other symptoms. They may pose a special health risk for infants, young children, some of the elderly, and people with severely compromised immune systems.
Total Coliform Bacteria (federal Revised Total Coliform Rule)			N/A	Naturally present in the environment	Coliforms are bacteria that are naturally present in the environment and are used as an indicator that other, potentially harmful, waterborne pathogens may be present or that a potential pathway exists through which contamination may enter the drinking water distribution system. We found coliforms indicating the need to look for potential problems in water treatment or distribution. When this occurs, we are required to conduct assessment(s) to identify problems and to correct any problems that were found during these assessments.

Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
E. coli (federal Revised Total Coliform Rule)		Footnote	(0)	Human and animal fecal waste	E. coli are bacteria whose presence indicates that the water may be contaminated with human or animal wastes. Human pathogens in these wastes can cause short-term effects, such as diarrhea, cramps, nausea, headaches, or other symptoms. They may pose a greater health risk for infants, young children, the elderly, and people with severely-compromised immune systems.
					For the consumer confidence report, if a water system detects E. coli and has violated the E. coli MCL, the water system shall include the following statements, as appropriate.
					 We had an <i>E. coli</i>-positive repeat sample following a total coliform-positive routine sample. We had a total coliform-positive repeat sample following an <i>E. coli</i>-coliform-positive repeat sample following an <i>E. colif-coliform-positive</i> repeat sample following and <i>E. colif-coliform-positive</i> repeat sample following an <i>E. colif-coliform-positive</i> repeat sample following and <i>E. colif-coliform-positive</i> repeat sample following repeat sample repe
					 We failed to take all required repeat samples following an <i>E. coli</i>-positive routine sample. We failed to test for <i>E. coli</i> when any re repeat sample tests positive for total coliform.
					If the E. coli MCL was not violated, the water system may include a statement that explains that although E. coli was detected, the water system is not in violation of the E. coli MCL.
<i>E. coli</i> (federal Revised Total Coliform Rule)		ŢŢ	N/A	Human and animal fecal waste	E. coli are bacteria whose presence indicates that the water may be contaminated with human or animal wastes. Human pathogens in these wastes can cause short-term effects, such as diarrhea, cramps, nausea, headaches, or other symptoms. They may pose a greater health risk for infants, young children, the elderly, and people with severely-compromised immune systems.
Fecal Indicator (E. coli) (Ground Water Rule)		0	(0)	Human and animal fecal waste	Fecal coliforms and <i>E. coli</i> are bacteria whose presence indicates that the water may be contaminated with human or animal wastes. Microbes in these wastes can cause short-term effects, such as diarrhea, cramps, nausea, headaches, or other symptoms. They may pose a special health risk for infants, young children, some of the elderly, and people with severely compromised immune systems.

¹ Routine and repeat samples are total coliform-positive and either is *E. coli*-positive or system fails to take repeat samples following *E. coli*-positive routine sample or system fails to analyze total coliform-positive repeat sample for *E. coli*.

Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Fecal Indicators (enterococci or coliphage) (Ground Water Rule)			N/A	Human and animal fecal waste	Fecal indicators are microbes whose presence indicates that the water may be contaminated with human or animal wastes. Microbes in these wastes can cause short-term effects, such as diarrhea, cramps, nausea, headaches, or other symptoms. They may pose a special health risk for infants, young children, some of the elderly, and people with severely compromised immune systems.
Turbidity		TT	N/A	Soil runoff	Turbidity has no health effects. However, high levels of turbidity can interfere with disinfection and provide a medium for microbial growth. Turbidity may indicate the presence of disease-causing organisms. These organisms include bacteria, viruses, and parasites that can cause symptoms such as nausea, cramps, diarrhea, and associated headaches.
Giardia lamblia, Viruses, Heterotrophic Plate Count Bacteria, Legionella, Cryptosporidium		TT	HPC = N/A; Others = (0)	Naturally present in the environment	Inadequately treated water may contain disease-causing organisms. These organisms include bacteria, viruses, and parasites that can cause symptoms such as nausea, cramps, diarrhea, and associated headaches.

Radioactive Contaminants

Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Gross Beta Particle Activity	pCi/L	50^{2}	(0)	Decay of natural and man-made deposits	Certain minerals are radioactive and may emit forms of radiation known as photons and beta radiation. Some people who drink water containing
					beta and photon emitters in excess of the MCL over many years may have an increased risk of getting cancer.
Strontium-90	pCi/L	8	0.35	Decay of natural and	Decay of natural and Some people who drink water containing strontium-90 in excess of the
Administra		***************************************		man-made deposit	MICE OVER MAILY YEARS MAY HAVE AN MICEASED HON OF SCHING CANCEL
Tritium	pCi/L	20,000	400	þ	Some people who drink water containing tritium in excess of the MCL
				man-made deposits	Over many years may have an increased his of securing cancer.
Gross Alpha Particle Activity	pCi/L	15	0	Erosion of natural	Certain minerals are radioactive and may emit a form of radiation
c				deposits	known as alpha radiation. Some people who drink water containing
					alpha emitters in excess of the MCL over many years may have an
					increased risk of getting cancer.

² Effective June 11, 2006, the gross beta particle activity MCL is 4 millirems/year annual dose equivalent to the total body or any internal organ. 50 pCi/L is used as a screening level.

Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Combined Radium 226 & 228	pCi/L	5	£(0) ³	Erosion of natural deposits	Some people who drink water containing radium 226 or 228 in excess of the MCL over many years may have an increased risk of getting cancer.
Total Radium (for nontransient-noncommunity water systems)	pCi/L	S	N/A	Erosion of natural deposits	Some people who drink water containing radium 223, 224, or 226 in excess of the MCL over many years may have an increased risk of getting cancer.
Uranium	pCi/L	20	0.43	Erosion of natural deposits	Some people who drink water containing uranium in excess of the MCL over many years may have kidney problems or an increased risk of getting cancer.

Inorganic Contaminants

Contaminant	Unit Measure -ment	MCL (AL) TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Aluminum	mg/L	-	9.0	Erosion of natural deposits; residue from some surface water treatment processes	Some people who drink water containing aluminum in excess of the MCL over many years may experience short-term gastrointestinal tract effects.
Antimony	T/Sri	9		Discharge from petroleum refineries; fire retardants; ceramics; electronics; solder	Some people who drink water containing antimony in excess of the MCL over many years may experience increases in blood cholesterol and decreases in blood cholesterol
Arsenic	ηg/L	10	0.004	Erosion of natural deposits; runoff from orchards; glass and electronics production wastes	Some people who drink water containing arsenic in excess of the MCL over many years may experience skin damage or circulatory system problems, and may have an increased risk of getting cancer.
Asbestos	MFL	7	7	Internal corrosion of asbestos cement water mains; erosion of natural deposits	Some people who drink water containing asbestos in excess of the MCL over many years may have an increased risk of developing benign intestinal polyns.
Barium	mg/L		2	Discharge of oil drilling wastes and from metal refineries; erosion of natural deposits	Some people who drink water containing barium in excess of the MCL over many years may experience an increase in blood pressure.
Beryllium	ng/L	4	***	Discharge from metal refineries, coal-burning factories, and electrical, aerospace, and defense industries	Some people who drink water containing beryllium in excess of the MCL over many years may develop intestinal lesions.

³ If reporting results for Ra-226 and Ra-228 as individual constituents, the PHG is 0.05 pCi/L for Ra-226 and 0.019 pCi/L for Ra-228.

					L. L
Contaminant	Unit Measure -ment	MCL (AL) TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Cadmium	T/Sn	\$	0.04	Internal corrosion of galvanized pipes; erosion of natural deposits; discharge from electroplating and industrial chemical factories, and metal refineries; runoff from waste batteries and paints	Some people who drink water containing cadmium in excess of the MCL over many years may experience kidney damage.
Chromium (Total)	µg/L	50	(100)	Discharge from steel and pulp mills and chrome plating; erosion of natural deposits	Some people who use water containing chromium in excess of the MCL over many years may experience allergic dermatitis.
Copper	mg/L	(AL=1.3)	0.3	Internal corrosion of household plumbing systems; erosion of natural deposits; leaching from wood preservatives	Copper is an essential nutrient, but some people who drink water containing copper in excess of the action level over a relatively short amount of time may experience gastrointestinal distress. Some people who drink water containing copper in excess of the action level over many years may suffer liver or kidney damage. People with Wilson's Disease should consult their personal doctor.
Cyanide	µg/L	150	150	Discharge from steel/metal, plastic and fertilizer factories	Some people who drink water containing cyanide in excess of the MCL over many years may experience nerve damage or thyroid problems.
Fluoride	mg/L	2.0	1	Erosion of natural deposits; water additive which promotes strong teeth; discharge from fertilizer and aluminum factories	Some people who drink water containing fluoride in excess of the federal MCL of 4 mg/L over many years may get bone disease, including pain and tenderness of the bones. Children who drink water containing fluoride in excess of the state MCL of 2 mg/L may get mottled teeth.
Lead	Л /8́п	(AL=15)	0.2	Internal corrosion of household water plumbing systems; discharges from industrial manufacturers; erosion of natural deposits	Infants and children who drink water containing lead in excess of the action level may experience delays in their physical or mental development. Children may show slight deficits in attention span and learning abilities. Adults who drink this water over many years may develop kidney problems or high blood pressure.
Mercury (Inorganic)	µg/L	2	1.2	Erosion of natural deposits; discharge from refineries and factories; runoff from landfills and cropland	Some people who drink water containing mercury in excess of the MCL over many years may experience mental disturbances, or impaired physical coordination, speech and hearing.
Nickel	µg/L	100	1.2	Erosion of natural deposits; discharge from metal factories	Some people who drink water containing nickel in excess of the MCL over many years may experience liver and heart effects.

Contaminant	Unit Measure -ment	MCL (AL) TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Nitrate (as Nitrogen, N)	mg/L	10	10	Runoff and leaching from fertilizer use; leaching from septic tanks and sewage; erosion of natural deposits	Infants below the age of six months who drink water containing nitrate in excess of the MCL may quickly become seriously ill and, if untreated, may die because high nitrate levels can interfere with the capacity of the infant's blood to carry oxygen. Symptoms include shortness of breath and blueness of the skin. High nitrate levels may also affect the oxygen-carrying ability of the blood of pregnant women.
Nitrite (as nitrogen, N)	mg/L	1		Runoff and leaching from fertilizer use; leaching from septic tanks and sewage; erosion of natural deposits	Infants below the age of six months who drink water containing nitrite in excess of the MCL may quickly become seriously ill and, if untreated, may die. Symptoms include shortness of breath and blueness of the skin.
Perchlorate	µg/L	9	_	Perchlorate is an inorganic chemical used in solid rocket propellant, fireworks, explosives, flares, matches, and a variety of industries. It usually gets into drinking water as a result of environmental contamination from historic aerospace or other industrial operations that used or use, store, or dispose of perchlorate and its salts.	Perchlorate has been shown to interfere with uptake of iodide by the thyroid gland, and to thereby reduce the production of thyroid hormones, leading to adverse affects associated with inadequate hormone levels. Thyroid hormones are needed for normal prenatal growth and development of the fetus, as well as for normal growth and development in the infant and child. In adults, thyroid hormones are needed for normal metabolism and mental function.
Selenium	µg/L	50	30	Discharge from petroleum, glass, and metal refineries; erosion of natural deposits; discharge from mines and chemical manufacturers; runoff from livestock lots (feed additive)	Selenium is an essential nutrient. However, some people who drink water containing selenium in excess of the MCL over many years may experience hair or fingernail losses, numbness in fingers or toes, or circulation system problems.
Thallium	Л/в́п	2	0.1	Leaching from ore-processing sites; discharge from electronics, glass, and drug factories	Some people who drink water containing thallium in excess of the MCL over many years may experience hair loss, changes in their blood, or kidney, intestinal, or liver problems.

Synthetic Organic Contaminants including Pesticides and Herbicides

Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
2,4-D	T/Sri	70	20	Runoff from herbicide used on row crops, range land, lawns, and aquatic weeds	Some people who use water containing the weed killer 2,4-D in excess of the MCL over many years may experience kidney, liver, or adrenal gland problems.
2,4,5-TP (Silvex)	J/gn	50	m	Residue of banned herbicide	Some people who drink water containing Silvex in excess of the MCL over many years may experience liver problems.
Acrylamide		T	(0)	Added to water during sewage/wastewater treatment	Some people who drink water containing high levels of acrylamide over a long period of time may experience nervous system or blood problems, and may have an increased risk of getting cancer.
Alachlor	T/gu	C1	4	Runoff from herbicide used on row crops	Some people who use water containing alachlor in excess of the MCL over many years may experience eye, liver, kidney, or spleen problems, or experience anemia, and may have an increased risk of getting cancer.
Atrazine	T/gri	ш	0.15	Runoff from herbicide used on row crops and along railroad and highway right-of-ways	Some people who use water containing atrazine in excess of the MCL over many years may experience cardiovascular system problems or reproductive difficulties.
Bentazon	T/Sri	18	200	Runoff/leaching from herbicide used on beans, peppers, corn, peanuts, rice, and ornamental grasses	Some people who drink water containing bentazon in excess of the MCL over many year may experience prostate and gastrointestinal effects.
Benzo(a)pyrene (PAH)	ng/L	200	7	Leaching from linings of water storage tanks and distribution mains	Some people who use water containing benzo(a)pyrene in excess of the MCL over many years may experience reproductive difficulties and may have an increased risk of getting cancer.
Carbofuran	T/Srl	18	0.7	Leaching of soil fumigant used on rice and alfalfa, and grape vineyards	Some people who use water containing carbofuran in excess of the MCL over many years may experience problems with their blood, or nervous or reproductive system problems.
Chlordane	Л/gu	100	30	Residue of banned insecticide	Some people who use water containing chlordane in excess of the MCL over many years may experience liver or nervous system problems, and may have an increased risk of getting cancer.
Dalapon	7/6п	200	790	Runoff from herbicide used on rights-of-ways, and crops and landscape maintenance	Some people who drink water containing dalapon in excess of the MCL over many years may experience minor kidney changes.

Contaminant	Unit Measure	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Di(2-ethylhexyl) adipate	ng/L	400	200	Discharge from chemical factories	Some people who drink water containing di(2-ethylhexyl) adipate in excess of the MCL over many years may experience weight loss, liver enlargement, or possible reproductive difficulties.
Di(2-ethylhexyl) phthalate	T/8ri	4	13	Discharge from rubber and chemical factories; inert ingredient in pesticides	Some people who use water containing di(2-ethylhexyl) phthalate in excess of the MCL over many years may experience liver problems or reproductive difficulties, and may have an increased risk of getting cancer.
Dibromochloropropane (DBCP)	ng/L	200	1.7	Banned nematocide that may still be present in solls due to runoff/leaching from former use on soybeans, cotton, vineyards, tomatoes, and tree fruit	Some people who use water containing DBCP in excess of the MCL over many years may experience reproductive difficulties and may have an increased risk of getting cancer.
Dinoseb	µg/L	7	14	Runoff from herbicide used on soybeans, vegetables, and fruits	Some people who drink water containing dinoseb in excess of the MCL over many years may experience reproductive difficulties.
Dioxin (2,3,7,8-TCDD)	Dg/L	30	0.05	Emissions from waste incineration and other combustion; discharge from chemical factories	Some people who use water containing dioxin in excess of the MCL over many years may experience reproductive difficulties and may have an increased risk of getting cancer.
Diquat	ng/L	20	9	Runoff from herbicide use for terrestrial and aquatic weeds	Some people who drink water containing diquat in excess of the MCL over many years may get cataracts.
Endothall	ug/L	100	94	Runoff from herbicide use for terrestrial and aquatic weeds; defoliant	Some people who drink water containing endothall in excess of the MCL over many years may experience stomach or intestinal problems.
Endrin	J/gri	2	0.3	Residue of banned insecticide and rodenticide	Some people who drink water containing endrin in excess of the MCL over many years may experience liver problems.
Epichlorohydrin		ŢŢ	(0)	Discharge from industrial chemical factories; impurity of some water treatment chemicals	Some people who drink water containing high levels of epichlorohydrin over a long period of time may experience stomach problems, and may have an increased risk of getting cancer.
Ethylene dibromide (EDB)	T/gn	50	10	Discharge from petroleum refineries; underground gas tank leaks; banned nematocide that may still be present in soils due to runoff and leaching from grain and fruit crops	Some people who use water containing ethylene dibromide in excess of the MCL over many years may experience liver, stomach, reproductive system, or kidney problems, and may have an increased risk of getting cancer.

Contaminant	Unit Measure	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Glyphosate	T/Bri	700	006	Runoff from herbicide use	Some people who drink water containing glyphosate in excess of the MCL over many years may experience kidney problems or reproductive difficulties.
Heptachlor	T/gn	10	8	Residue of banned insecticide	Some people who use water containing heptachlor in excess of the MCL over many years may experience liver damage and may have an increased risk of getting cancer.
Heptachlor epoxide	T/Su	10	9	Breakdown of heptachlor	Some people who use water containing heptachlor epoxide in excess of the MCL over many years may experience liver damage, and may have an increased risk of getting cancer.
Hexachlorobenzene	µg/L		0.03	Discharge from metal refineries and agricultural chemical factories; byproduct of chlorination reactions in wastewater	Some people who drink water containing hexachlorobenzene in excess of the MCL over many years may experience liver or kidney problems, or adverse reproductive effects, and may have an increased risk of getting cancer.
Hexachlorocyclopentadiene	T/8n	50	2	Discharge from chemical factories	Some people who use water containing hexachlorocyclopentadiene in excess of the MCL over many years may experience kidney or stomach problems.
Lindane	ng/L	200	32	Runoff/leaching from insecticide used on cattle, lumber, and gardens	Some people who drink water containing lindane in excess of the MCL over many years may experience kidney or liver problems.
Methoxychlor	T/Bri	30	60:0	Runoff/leaching from insecticide used on fruits, vegetables, alfalfa, and livestock	Some people who drink water containing methoxychlor in excess of the MCL over many years may experience reproductive difficulties.
Molinate (Ordram)	T/Bri	20		Runoff/leaching from herbicide used on rice	Some people who use water containing molinate in excess of the MCL over many years may experience reproductive effects.
Oxamyl (Vydate)	hg/L	50	26	Runoff/leaching from insecticide used on field crops, fruits and ornamentals, especially apples, potatoes, and tomatoes	Some people who drink water containing oxamyl in excess of the MCL over many years may experience slight nervous system effects.
PCBs (Polychlorinated biphenyls)	J/gn	500	06	Runoff from landfills; discharge of waste chemicals	Some people who drink water containing PCBs in excess of the MCL over many years may experience changes in their skin, thymus gland problems, immune deficiencies, or reproductive or nervous system difficulties, and may have an increased risk of getting cancer.
Pentachlorophenol	T/SH	-	0.3	Discharge from wood preserving factories, cotton and other insecticidal/herbicidal uses	Some people who use water containing pentachlorophenol in excess of the MCL over many years may experience liver or kidney problems, and may have an increased risk of getting cancer.

		-			
Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Picloram	T/Sri	200	166	Herbicide runoff	Some people who drink water containing picloram in excess of the MCL over many years may experience liver problems.
Simazine	T/Sri	4	4	Herbicide runoff	Some people who use water containing simazine in excess of the MCL over many years may experience blood problems.
Thiobencarb	T/Sri	70	42	Runoff/leaching from herbicide used on rice	Some people who use water containing thiobencarb in excess of the MCL over many years may experience body weight and blood effects.
Toxaphene	Lg/L	rn	0.03	Runoff/leaching from insecticide used on cotton and cattle	Some people who use water containing toxaphene in excess of the MCL over many years may experience kidney, liver, or thyroid problems, and may have an increased risk of getting cancer.
1,2,3-Trichloropropane	ng/L	\$	0.7	Discharge from industrial and agricultural chemical factories; leaching from hazardous waste sites; used as cleaning and maintenance solvent, paint and varnish remover, and cleaning and degreasing agent; byproduct during the production of other compounds and pesticides.	Some people who drink water containing 1,2,3- trichloropropane in excess of the MCL over many years may have an increased risk of getting cancer.

Volatile Organic Contaminants

Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Вепzепе	µg/L	1	0.15	Discharge from plastics, dyes and nylon factories; leaching from gas storage tanks and landfills	Some people who use water containing benzene in excess of the MCL over many years may experience anemia or a decrease in blood platelets, and may have an increased risk of getting cancer.
Carbon tetrachloride	ng/L	500	100	Discharge from chemical plants and other industrial activities	Some people who use water containing carbon tetrachloride in excess of the MCL over many years may experience liver problems and may have an increased risk of getting cancer.
1,2-Dichlorobenzene	T/Sn	009	009	Discharge from industrial chemical factories	Some people who drink water containing 1,2-dichlorobenzene in excess of the MCL over many years may experience liver, kidney, or circulatory system problems.

Contaminant	Unit Measure	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
1,4-Dichlorobenzene	T/Sin	S	9	Discharge from industrial chemical factories	Some people who use water containing 1.4-dichlorobenzene in excess of the MCL over many years may experience anemia, liver, kidney, or spleen damage, or changes in their blood.
1,1-Dichloroethane	7/8n	S	'n	Extraction and degreasing solvent; used in the manufacture of pharmaceuticals, stone, clay, and glass products; fumigant	Some people who use water containing 1,1-dichloroethane in excess of the MCL over many years may experience nervous system or respiratory problems.
1,2-Dichloroethane	ng/L	200	400	Discharge from industrial chemical factories	Some people who use water containing 1,2- dichloroethane in excess of the MCL over many years may have an increased risk of getting cancer.
1,1-Dichloroethylene	hg/L	9	10	Discharge from industrial chemical factories	Some people who use water containing 1,1- dichloroethylene in excess of the MCL over many years may experience liver problems.
cis-1,2-Dichloroethylene	7/8п	9	100	Discharge from industrial chemical factories; major biodegradation byproduct of TCE and PCE groundwater contamination	Some people who use water containing cis-1,2-dichloroethylene in excess of the MCL over many years may experience liver problems.
trans-1,2-Dichloroethylene	T/Str	10	09	Discharge from industrial chemical factories; minor biodegradation byproduct of TCE and PCE groundwater contamination	Some people who drink water containing trans-1,2-dichloroethylene in excess of the MCL over many years may experience liver problems.
Dichloromethane	T/Sn	v	4	Discharge from pharmaceutical and chemical factories; insecticide	Some people who drink water containing dichloromethane in excess of the MCL over many years may experience liver problems and may have an increased risk of getting cancer.
1,2-Dichloropropane	T/Stl	5	0.5	Discharge from industrial chemical factories; primary component of some fumigants	Some people who use water containing 1,2-uromorphopano in excess of the MCL over many years may have an increased risk of getting cancer.
1,3-Dichloropropene	ng/L	500	200	Runoff/leaching from nematocide used on croplands	Some people who use water containing 1,5-dichloropropene in excess of the MCL over many years may have an increased risk of getting cancer.
Ethylbenzene	Л/Sri	300	300	Discharge from petroleum refineries; industrial chemical factories	excess of the MCL over many years may experience liver or kidney problems.
Methyl-tert-butyl ether	hg/L		13	Leaking underground storage tanks; discharges from petroleum and chemical factories	Some people who use water containing memorizer rough ether in excess of the MCL over many years may have an increased risk of getting cancer.

	Unit	MCL	PHG	Major Sources of	The 1st of the American
Contaminant	Measure -ment	TT, as noted	(MCLG)	Contamination	Leann Ellects Language
Monochlorobenzene	ng/L	70	70	Discharge from industrial and agricultural chemical factories	Some people who use water containing monochlorobenzene in excess of the MCL over many years may experience liver or kidney problems
Styrene	µg/L	100	0.5	Discharge from rubber and plastic factories; leaching from landfills	Some people who drink water containing styrene in excess of the MCL over many years may experience liver, kidney, or circulatory system problems, and may have an increased risk of getting cancer.
1,1,2,2-Tetrachloroethane	J/gµ		0.1	Discharge from industrial and agricultural chemical factories; solvent used in production of TCE, pesticides, varnish and lacquers	Some people who drink water containing 1,1,2,2-tetrachloroethane in excess of the MCL over many years may experience liver or nervous system problems.
Tetrachloroethylene (PCE)	µg/L	\$	90.0	Discharge from factories, dry cleaners, and auto shops (metal degreaser)	Some people who use water containing tetrachloroethylene in excess of the MCL over many years may experience liver problems, and may have an increased risk of getting cancer.
1,2,4-Trichlorobenzene	ng/L	5	3	Discharge from textile-finishing factories	Some people who use water containing 1,2,4-trichlorobenzene in excess of the MCL over many years may experience adrenal gland changes.
1,1,1-Trichloroethane	µg/L	200	1000	Discharge from metal degreasing sites and other factories; manufacture of food wrappings	Some people who use water containing 1,1,1-trichloroethane in excess of the MCL over many years may experience liver, nervous system, or circulatory system problems.
1,1,2-Trichloroethane	ng/L	5	0.3	Discharge from industrial chemical factories	Some people who use water containing 1,1,2-trichloroethane in excess of the MCL over many years may experience liver, kidney, or immune system problems.
Trichloroethylene (TCE)	T/Sn	5	1.7	Discharge from metal degreasing sites and other factories	Some people who use water containing trichloroethylene in excess of the MCL over many years may experience liver problems and may have an increased risk of getting cancer.
Toluene	T/8n	150	150	Discharge from petroleum and chemical factories; underground gas tank leaks	Some people who use water containing toluene in excess of the MCL over many years may experience nervous system, kidney, or liver problems.
Trichlorofluoromethane	T/Sri	150	1300	Discharge from industrial factories; degreasing solvent; propellant and refrigerant	Some people who use water containing trichlorofluoromethane in excess of the MCL over many years may experience liver problems.
1,1,2-Trichloro-1,2,2- trifluoroethane	mg/L	1.2	4	Discharge from metal degreasing sites and other factories; drycleaning solvent; refrigerant	Some people who use water containing 1,1,2-trichloro-1,2,2-trifloroethane in excess of the MCL over many years may experience liver problems.

Contaminant	Unit Measure -ment	MCL TT, as noted	PHG (MCLG)	Major Sources of Contamination	Health Effects Language
Vinyl chloride	T/gu	500	50	Leaching from PVC piping; discharge from plastics factories; biodegradation byproduct of TCE and PCE groundwater contamination	Some people who use water containing vinyl chloride in excess of the MCL over many years may have an increased risk of getting cancer.
Xylenes	mg/L	1.750	1.8	Discharge from petroleum and chemical factories; fuel solvent	Some people who use water containing xylenes in excess of the MCL over many years may experience nervous system damage.

Disinfection Byproducts, Disinfectant Residuals, and Disinfection Byproduct Precursors

MCL PHG [MRDL] (MCLG) IT, as noted [MRDLG]
N/A Byproduct of drinking water disinfection
N/A Byproduct of drinking water disinfection
0.1 Byproduct of drinking water disinfection
[MRDLG = Drinking water disinfectant 4 (as Cl ₂₎] added for treatment
[MRDLG = Drinking water disinfectant 4 (as Cl ₂) added for treatment

	TINE	MOT	SHO	TOWN THE TAX TO THE TA	- потпитентический потп
11000				Major Sources of	A COUNTY AND
Contaminant	Measure -ment	[MKDL] TT, as noted	(MCLG) [MRDLG]	Contamination	Health Effects Language
Chlorite	T/8m	1.0	0.05	Byproduct of drinking water	Some infants and young children who drink water
				disinfection	containing chlorite in excess of the MCL could experience
					nervous system effects. Similar effects may occur in
					fetuses of pregnant women who drink water containing
					chlorite in excess of the MCL. Some people may
					experience anemia.
Chlorine Dioxide	T/gn	[MRDL =	[MRDLG =	Drinking water disinfectant	Some infants and young children who drink water
		800 (as CIO ₂)]	800 (as	added for treatment	containing chlorine dioxide in excess of the MRDL could
			$ClO_2)]$		experience nervous system effects. Similar effects may
					occur in fetuses of pregnant women who drink water
					containing chlorine dioxide in excess of the MRDL. Some
					people may experience anemia.
Control of DBP precursors		TT	N/A	Various natural and man-made	Total organic carbon (TOC) has no health effects.
(TOC)				sources	However, total organic carbon provides a medium for the
					formation of disinfection byproducts. These byproducts
					include trihalomethanes (THMs) and haloacetic acids
					(HAAs). Drinking water containing these byproducts in
					excess of the MCL may lead to adverse health effects, liver
					or kidney problems, or nervous system effects, and may
					lead to an increased risk of cancer.

APPENDIX B: Regulated Contaminants with Secondary Drinking Water Standards

Monitoring Required by Section 64449, Chapter 15, Title 22, California Code of Regulations

Contaminant	Unit Measurement	MCL	Typical Source of Contaminant
Aluminum	μg/L	200	Erosion of natural deposits; residual from some surface water treatment processes
Color	Units	15	Naturally-occurring organic materials
Copper	mg/L	1.0	Internal corrosion of household plumbing systems; erosion of natural deposits; leaching from wood preservatives
Foaming Agents (MBAS)	μg/L	500	Municipal and industrial waste discharges
Iron	μg/L	300	Leaching from natural deposits; industrial wastes
Manganese	μg/L	50	Leaching from natural deposits
Methyl-tert-butyl ether (MTBE)	μg/L	5	Leaking underground storage tanks; discharge from petroleum and chemical factories
OdorThreshold	Units	3	Naturally-occurring organic materials
Silver	μg/L	100	Industrial discharges
Thiobencarb	μg/L	1	Runoff/leaching from rice herbicide
Turbidity	Units	5	Soil runoff
Zinc	mg/L	5.0	Runoff/leaching from natural deposits; industrial wastes
Total Dissolved Solids (TDS)	mg/L	1,000	Runoff/leaching from natural deposits
Specific Conductance	μS/cm	1,600	Substances that form ions when in water; seawater influence
Chloride	mg/L	500	Runoff/leaching from natural deposits; seawater influence
Sulfate	mg/L	500	Runoff/leaching from natural deposits; industrial wastes

There are no PHGs, MCLGs, or mandatory standard health effects language for these constituents because secondary MCLs are set on the basis of aesthetics.

APPENDIX C: Monitored Contaminants with No MCLs

Background

The 1996 Amendments to the SDWA required the U.S. EPA to establish criteria for a monitoring program for unregulated contaminants, and to publish, once every five years, a list of no more than 30 contaminants to be monitored by public water systems (PWS).

Section 64450 of the California Code of Regulations also required certain water systems to monitor a number of unregulated contaminants, with contaminant lists that were published or revised in 1990, 1996, 2000, and 2003. This section of the California Code of Regulations was repealed effective October 18, 2007. Water systems that continued to monitor for state unregulated contaminants are encouraged, but not required, to include the information regarding detected contaminants in the CCR.

Although Section 64450 of the California Code of Regulations was repealed, the State Water Board may request water systems to monitor for specific contaminants per HSC section 116375(b).

Federal UCMR 1 (2001 – 2003 Monitoring)

The U.S. EPA published the first list of contaminants to monitor as part of the UCMR in September 1999. Contaminants were divided into two lists: Assessment Monitoring (List 1), and Screening Survey (List 2).

Assessment Monitoring of List 1 contaminants was conducted by large PWS serving more than 10,000 people and 800 representative small PWS serving 10,000 or fewer people. Assessment Monitoring was conducted by each PWS over a 12-month period between 2001 and 2003.

Screening Survey was conducted by a randomly selected set of 300 large and small PWSs for List 2 contaminants. Screening Survey for chemical contaminants was conducted in 2001 and 2002 for small and large PWS, respectively. Screening Survey for *Aeromonas* was conducted in 2003 for small and large PWS.

	UCMR 1	
List 1 – Assessment Monitoring	List 2 – Screening Survey	
2,4-dinitrotoluene	1,2-diphenylhydrazine	
2,6-dinitrotoluene	2-methyl-phenol	
Acetochlor	2,4-dichlorophenol	
DCPA mono-acid degradate	2,4-dinitrophenol	
DCPA di-acid degradate	2,4,6-trichlorophenol	
4,4'-DDE	Aeromonas	
EPTC	Alachlor ESA	
Molinate Diazinon		
MTBE	Disulfoton	
Nitrobenzene	Diuron	
Perchlorate	Fonofos	

Terbacil	Linuron
	Nitrobenzene
	Prometon
6	Hexahydro-1,3,5-trinitro-1-3-5-triazine [RDX]
	Terbufos

Federal UCMR 2 (2008 – 2010 Monitoring)

The U.S. EPA published the second list of contaminants to monitor as part of the UCMR in January 2007.

Assessment Monitoring was required of all PWS serving more than 10,000 people and 800 representative PWS serving 10,000 or fewer people for List 1 contaminants. Assessment Monitoring was required of each PWS during a 12-month period from January 2008 to December 2010.

Screening Survey was required of all PWS serving more than 100,000 people, 320 representative PWS serving 10,001 to 100,000 people, and 480 representative PWS serving 10,000 or fewer people for List 2 contaminants. Screening Survey was required of each PWS during a 12-month period from January 2008 to December 2010.

UC	CMR 2
List 1 – Assessment Monitoring	List 2 – Screening Survey
Dimethoate Terbufos sulfone 2,2',4,4'-tetrabromodiphenyl ether 2,2',4,4',5-pentabromodiphenyl ether 2,2',4,4',5,5'-hexabromodiphenyl 2,2',4,4',5,5'-hexabromodiphenyl ether 2,2',4,4',6-pentabromodiphenyl ether 1,3-dinitrobenzene 2,4,6-trinitrotoluene (TNT) Hexahydro-1,3,5-trinitro-1,3,5-trazine (RDX)	Acetochlor ethane sulfonic acid Acetochlor oxanilic acid Alachlor ethane sulfonic acid Alachlor oxanilic acid Metolachlor ethane sulfonic acid Metolachlor oxanilic acid Acetochlor Alachlor Metolachlor
	N-nitrosodiethylamine (NDEA) N-nitrosodimethylamine (NDMA) N-nitroso-di-n-butylamine (NDBA) N-nitroso-di-n-propylamine (NDPA) N-nitrosomethylethylamine (NMEA) N-nitrosopyrrolidine (NPYR)

Federal UCMR 3 (2013 – 2015 Monitoring)

The third UCMR list of contaminants was published in May 2012.

Assessment Monitoring (List 1 Contaminants) was required of all PWS serving more than 10,000 people and 800 representative PWS serving 10,000 or fewer people. Assessment

Monitoring was required of each PWS during a 12-month period from January 2013 to December 2015.

Screening Survey (List 2 Contaminants) was required of all PWS serving more than 100,000 people, 320 representative PWS serving 10,001 to 100,000 people, and 480 representative PWS serving 10,000 or fewer people. Screening Survey was required of each PWS during a 12-month period from January 2013 to December 2015.

Pre-screen Testing (List 3 Contaminants) was required from a selection of 800 representative PWS serving 1,000 or fewer people that do not disinfect. These PWS were selected because they have groundwater wells that were located in areas of karst or fractured bedrock. Monitored lasted 12 months between January 2013 and December 2015.

UCM	1R 3
List 1 – Assessment Monitoring	List 2 – Screening Survey
1,2,3-trichloropropane 1,3-butadiene Chloromethane (methyl chloride) 1,2-dichloroethane Bromomethane (methyl bromide) Chlorodifluoromethane (HCFC-22) Bromochloromethane (halon 1011) 1,4-dioxane	17-β-estradiol 17-α-ethynylestradiol (ethinyl estradiol) 16-α-hydroxyestradiol (estriol) Equilin Estrone Testosterone 4-anderostene-3,17-dione
Vanadium Molybdenum Cobalt Strontium Chromium (total) Chromium-6	List 3 – Pre-Screen Testing Enteroviruses Noroviruses
Perfluorooctanesulfonate acid (PFOS) Perfluorooctanoic acid (PFOA) Perfluorononanoic acid (PFNA) Perfluorohexanesulfonic acid (PFHxS) Perfluoroheptanoic acid (PFHpA) Perfluorobutanesulfonic acid (PFBS)	

Federal UCMR 4 (2018 – 2020 Monitoring)

The fourth list of contaminants to monitor as part of the UCMR was published by the U.S. EPA in December 2016.

PWSs are required to monitor for 10 cyanotoxins at the entry point to the distribution system during a 4-consecutive month period from March 2018 through November 2020, according to the table

below. PWSs are also required to monitor for 20 additional chemical contaminants and indicators during a 12-month period from January 2018 through December 2020. The sampling site for these additional chemicals is the entry point to the distribution system, except for HAAs that need to be monitored at the Stage 2 D/DBPR sampling sites. The two indicators, *i.e.*, TOC and bromide, need to be monitored at source water intakes.

System Size (Population Served)	10 Cyanotoxins	20 Chemicals
Small Systems (25 – 10,000)	800 randomly selected surface water or ground water under the direct influence of surface water (GWUDI) systems	A different group of 800 randomly selected surface water systems, GWUDI and groundwater systems
Large Systems (10,001 or more)	All surface water and GWUDI systems	All surface water, groundwater and GWUDI systems

The 10 cyanotoxins and 20 additional chemical contaminants and indicators are listed in the table below.

UCMR 4			
Cyanotoxins	Minimum Reporting Level		
Total Microcystin	0.3 μg/L		
Microcystin-LA	$0.008 \mu\mathrm{g/L}$		
Microcystin-LF	0.006 μg/L		
Microcystin-LR	0.02 μg/L		
Microcystin-LY	0.009 μg/L		
Microcystin-RR	0.006 μg/L		
Microcystin-YR	0.02 μg/L		
Nodularin	$0.005 \mu\mathrm{g/L}$		
Anatoxin-a	0.03 μg/L		
Cylindrospermopsin	0.09 µg/L		
Additional Chemicals	Minimum Reporting Level		
Germanium	0.3 μg/L		
Manganese	0.4 μg/L		
Alpha-hexachlorocyclohexane	0.01 μg/L		
Chlorpyrifos	0.03 μg/L		
Dimethipin	0.2 µg/L		
Ethoprop	0.03 μg/L		
Oxyfluorfen	0.05 μg/L		
Profenofos	0.3 μg/L		
Tebuconazole	0.2 μg/L		
Total Permethrin (cis- & trans-)	0.04 μg/L		
Tribufos	0.07 μg/L		
HAA5	N/A		
HAA6Br ¹	N/A		

HAA9 ²	N/A
1-butanol 2-methoxyethanol 2-propen-1-ol	2.0 μg/L 0.4 μg/L 0.5 μg/L
butylated hydroxyanisole	0.03 μg/L
o-toluidine	0.007 μg/L
quinoline	0.02 μg/L
Total Organic Carbon (TOC)	N/A
Bromide	N/A

¹ HAA6Br: Bromochloroacetic acid, bromodichloroacetic acid, dibromoacetic acid, dibromochloroacetic acid, monobromoacetic acid, and tribromoacetic acid.

Reporting

U.S. EPA is essentially silent on the issue of reporting federal UCMR contaminants beyond the previous calendar year's detections, other than to say it is not required and that data older than five years need not be reported. As a result, the State Water Board recommends systems to report data for five years from the date of the last sampling.

² HAA9: Bromochloroacetic acid, bromodichloroacetic acid, chlorodibromoacetic acid, dibromoacetic acid, dichloroacetic acid, monobromoacetic acid, monochloroacetic acid, tribromoacetic acid, and trichloroacetic acid.

APPENDIX D: State Contaminants with Notification Levels

Inclusion of the Notification Level (NL) and health effects language for contaminant concentrations detected above the NL is recommended, but not required.

Chemical	Notification Level	Health Effects Language (Optional)
Boron	1 mg/L	Boron exposures resulted in decreased fetal weight (developmental effects) in newborn rats.
n-Butylbenzene	260 μg/L	Exposures to cumene (isopropylbenzene), a surrogate for
sec-Butylbenzene	260 μg/L	n-, sec-, and tert-butylbenzene, resulted in increased
tert-Butylbenzene	260 μg/L	kidney weight in rats.
Carbon Disulfide	160 μg/L	Carbon disulfide exposures resulted in decreased motor conduction velocity in people.
Chlorate	800 μg/L	Animal studies demonstrated that chlorate exposure in rats caused adverse effects to the pituitary and thyroid glands.
2-Chlorotoluene	140 μg/L	2-Chlorotoluene exposures resulted in decrease in body
4-Chlorotoluene	140 μg/L	weight gain in rats. 4-Chlorotoluene is expected to have health effects similar to those of 2-chlorotoluene.
Diazinon	1.2 μg/L	Diazinon exposures may result in neurotoxic effects.
Dichlodifluoromethane [Freon 12]	1 mg/L	Dichlorodifluoromethane exposures resulted in reduced body weight in rats.
1,4-Dioxane	1 μg/L	1,4-Dioxane exposures resulted in cancer, based on studies in laboratory animals.
Ethylene Glycol	14 mg/L	Ethylene glycol exposures resulted in kidney toxicity in rats.
Formaldehyde	100 μg/L	Formaldehyde exposures resulted in reduced weight gain and histopathology in rats.
Octahydro-1,3,5,7- tetranitro-1,3,5,7- tetrazocine [HMX]	350 μg/L	Octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine exposures resulted in liver lesions in rats.
Isopropylbenzene	770 μg/L	Isopropylbenzene exposures resulted in increased kidney weight in rats.
Manganese	500 μg/L	Manganese exposures resulted in neurological effects. High levels of manganese in people have been shown to result in adverse effects to the nervous system.
Methyl Isobutyl Ketone [MIBK]	120 μg/L	Methyl isobutyl ketone exposures resulted in increased kidney and liver weight, and kidney pathology in rats.
Naphthalene	17 μg/L	Naphthalene exposures resulted in decreased body weight in rats.
N-Nitrosodiethylamine [NDEA]	10 ng/L	N-nitrosodiethylamine exposures resulted in cancer in a variety of laboratory animals.

Chemical	Notification Level	Health Effects Language (Optional)
N-Nitrosodimethylamine [NDMA]	10 ng/L	N-nitrosodimethylamine exposures resulted in cancer in a variety of laboratory animals.
N-Nitrosodi-n- propylamine [NDPA]	10 ng/L	N-nitrosodi-n-propylamine exposures resulted in cancer in a variety of laboratory animals.
Perfluorooctanoic Acid [PFOA]	5.1 ng/L**	Perfluorooctanoic acid exposures resulted in increased liver weight in laboratory animals.
Perfluorooctanesulfonic Acid [PFOS]	6.5 ng/L**	Perfluorooctanesulfonic acid exposures resulted in immune suppression, specifically, a decrease in antibody response to an exogenous antigen challenge.
Propachlor	90 μg/L	Propachlor exposures resulted in decrease in weight gain, decrease in food intake, and relative liver weight increase in rats.
n-Propylbenzene	260 μg/L	Exposures to cumene (isopropylene), a surrogate for n-propylbenzene, resulted in increased kidney weight in rats.
Hexahydro-1,3,5-trinitro- 1-3-5-triazine [RDX]	300 ng/L	Hexahydro-1,3,5-trinitro-1-3-5-triazine exposures resulted in liver carcinomas and adenomas in female mice.
Tertiary Butyl Alcohol [TBA]	12 μg/L	Tert-butyl alcohol exposures resulted in cancer in laboratory animals.
1,2,4-Trimethylbenzene	330 μg/L	1,2,4-Trimethylbenzene exposures resulted in increased serum phosphorus levels in rats.
1,3,5-Trimethylbenzene	330 μg/L	1,3,5-Trimethylbenzene exposures resulted in increased serum phosphorus levels in rats.
2,4,6-Trinitrotoluene [TNT]	l μg/L	2,4,6-Trinitrotoluene exposures resulted in urinary bladder transitional cell papillomas and squamous cell carcinomas in female rats.
Vanadium	50 μg/L	Vanadium exposures resulted in developmental and reproductive effects in rats.

^{**} The July 2018 notification levels for PFOA of 14 ng/L and PFOS of 13 ng/L were superseded on August 22, 2019 by new notification levels of 5.1 ng/L for PFOA and 6.5 ng/L for PFOS.

APPENDIX E: Special Language for Nitrate, Arsenic, Lead, Radon, *Cryptosporidium*, Ground Water Systems, and Surface Water Systems

(A) Nitrate: For systems that detect nitrate above 5 mg/L as nitrogen, but below 10 mg/L as nitrogen, the following language is REQUIRED:

Nitrate in drinking water at levels above 10 mg/L is a health risk for infants of less than six months of age. Such nitrate levels in drinking water can interfere with the capacity of the infant's blood to carry oxygen, resulting in serious illness; symptoms include shortness of breath and blueness of the skin. Nitrate levels above 10 mg/L may also affect the ability of the blood to carry oxygen in other individuals, such as pregnant women and those with specific enzyme deficiencies. If you are caring for an infant, or you are pregnant, you should ask advice from your health care provider.

If a utility cannot demonstrate to the State Water Board with at least five years of the most current monitoring data that its nitrate levels are stable, it must also add the following language to the preceding statement on nitrate:

Nitrate levels may rise quickly for short periods of time because of rainfall or agricultural activity.

(B) Arsenic: For systems that detect arsenic above 5 μ g/L, but below or equal to 10 μ g/L, the following language is REQUIRED:

While your drinking water meets the federal and state standard for arsenic, it does contain low levels of arsenic. The arsenic standard balances the current understanding of arsenic's possible health effects against the cost of removing arsenic from drinking water. The U.S. Environmental Protection Agency continues to research the health effects of low levels of arsenic, which is a mineral known to cause cancer in humans at high concentrations and is linked to other health effects such as skin damage and circulatory problems.

(C) Lead¹: Consistent with 40 CFR section 141.154(d)(1), every Consumer Confidence Report (CCR) must include the lead-specific language shown below. A water system may provide its own educational statement, but only after consulting with the State Water Board.

¹ All water systems are required to comply with the state Lead and Copper Rule (LCR). Water systems are also required to comply with the federal LCR, and its revisions and corrections. The 2007 Short-term Revisions of the LCR included mandatory language requirements that have not yet been adopted by the State Water Board.

If present, elevated levels of lead can cause serious health problems, especially for pregnant women and young children. Lead in drinking water is primarily from materials and components associated with service lines and home plumbing. [NAME OF UTILITY] is responsible for providing high quality drinking water, but cannot control the variety of materials used in plumbing components. When your water has been sitting for several hours, you can minimize the potential for lead exposure by flushing your tap for 30 seconds to 2 minutes before using water for drinking or cooking. [Optional: If you do so, you may wish to collect the flushed water and reuse it for another beneficial purpose, such as watering plants.] If you are concerned about lead in your water, you may wish to have your water tested. Information on lead in drinking water, testing methods, and steps you can take to minimize exposure is available from the Safe Drinking Water Hotline or at http://www.epa.gov/lead.

Consistent with the California Code of Regulations, section 64482(c), systems that detect lead above 15 μ g/L in more than 5 percent, and up to and including 10 percent, of sites sampled (or if your system samples fewer than 20 sites and has even one sample above the Action Level [AL]), the following language is REQUIRED:

Infants and young children are typically more vulnerable to lead in drinking water than the general population. It is possible that lead levels at your home may be higher than at other homes in the community as a result of materials used in your home's plumbing. If you are concerned about elevated lead levels in your home's water, you may wish to have your water tested and/or flush your tap for 30 seconds to 2 minutes before using tap water. Additional information is available from the U.S. EPA Safe Drinking Water Hotline (1-800-426-4791).

(D) Radon: Systems that performed monitoring that indicates the presence of radon in the finished water MUST include the results of the monitoring and an explanation of the significance of the results. The following language MAY be used:

We constantly monitor the water	supply for various	contaminants. We	have detected radon
in the finished water supply in	out of	samples tested.	There is no federal
regulation for radon levels in dr	inking water. Expo	osure over a long	period of time to air
transmitting radon may cause ad	dverse health effect	ts.	

The language below MAY be included if the level of information is helpful.

Radon is a radioactive gas that you cannot see, taste, or smell. It is found throughout the U.S. Radon can move up through the ground and into a home through cracks and holes in the foundation. Radon can build up to high levels in all types of homes. Radon can also get into indoor air when released from tap water from showering, washing dishes, and other household activities. Compared to radon entering the home through soil, radon entering the home through tap water will in most cases be a small source of radon in indoor air. Radon is a known human carcinogen. Breathing air containing radon can lead to lung cancer. Drinking water containing radon may also cause increased risk of

stomach cancer. If you are concerned about radon in your home, test the air in your home. Testing is inexpensive and easy. You should pursue radon removal for your home if the level of radon in your air is 4 picocuries per liter of air (pCi/L) or higher. There are simple ways to fix a radon problem that are not too costly. For additional information, call your State radon program (1-800-745-7236, the U.S. EPA Safe Drinking Water Act Hotline (1-800-426-4791), or the National Safe Council Radon Hotline (1-800-767-7236).

(E) Cryptosporidium: Systems that have performed any monitoring for Cryptosporidium that indicates that Cryptosporidium may be present in the source water or finished water MUST include the results of the monitoring and an explanation of the significance of the results. The following language MAY be used:

Cryptosporidium is a microbial pathogen found in surface water throughout the U.S. Although filtration removes Cryptosporidium, the most commonly-used filtration methods cannot guarantee 100 percent removal. Our monitoring indicates the presence of these organisms in our source water and/or finished water. Current test methods do not allow us to determine if the organisms are dead or if they are capable of causing disease. Ingestion of Cryptosporidium may cause cryptosporidiosis, an abdominal infection. Symptoms of infection include nausea, diarrhea, and abdominal cramps. Most healthy individuals can overcome the disease within a few weeks. However, immuno-compromised people, infants, small children, and the elderly are at greater risk of developing lifethreatening illness. We encourage immuno-compromised individuals to consult their doctor regarding appropriate precautions to take to avoid infection. Cryptosporidium must be ingested to cause disease, and it may be spread through means other than drinking water.

(F) Groundwater Systems: For ground water systems that had a treatment technique (TT) violation described in Item S of the document titled "Instructions for Completing the 2018 CCR for Small Water Systems", the following language MAY be used to describe the potential health effects. The U.S. Environmental Protection Agency (EPA) did not provide standard health effect language for these TT violations in the Ground Water Rule; U.S. EPA provided the language in their guidance to water systems.

Inadequately protected or treated water may contain disease-causing organisms. These organisms can cause symptoms such as diarrhea, nausea, cramps, and associated headaches.

(G) Surface Water Systems: For surface water systems that had a TT violation under the Surface Water Treatment Rule (SWTR), Interim Enhanced Surface Water Treatment Rule (IESWTR), Filter Backwash Recycling Rule (FBRR), or Long-term 1 Enhanced Surface Water Treatment Rule (LT1ESWTR), as described in Item U of the document titled "Instructions for Completing the 2018 CCR for Small Water Systems", the following language is REQUIRED to describe the potential health effects:

Inadequately treated water may contain disease-causing organisms. These organisms include bacteria, viruses, and parasites that can cause symptoms such as nausea, cramps, diarrhea, and associated headaches.

For surface water systems that had a TT violation under the Long-term 2 Enhanced Surface Water Treatment Rule (LT2ESWTR), as described in Item U of the document titled "Instructions for Completing the 2018 CCR for Small Water Systems", the following language MAY be used to describe the potential health effects. U.S. EPA did not provide standard health effect language for these TT violations in the LT2ESWTR; U.S. EPA provided the language in their guidance to water systems.

LT2ESWTR TT Violation	Health Effect Language	
Uncovered and Untreated Finished Water Reservoir	Inadequately protected water may contain disease-causing organisms. These organisms can cause symptoms such as diarrhea, nausea, cramps, and associated headaches.	
Determine and Report Bin Classification	Inadequately treated water may contain disease-causing organisms. These organisms can cause symptoms such as diarrhea, nausea, cramps, and associated headaches.	
Provide or Install an Additional Level of Treatment	Inadequately treated water may contain disease-causing organisms. These organisms can cause symptoms such as diarrhea, nausea, cramps, and associated headaches.	

APPENDIX F: Certification Form

Consumer Confidence Report Certification Form

(to be submitted with a copy of the CCR)

(To certify electronic delivery of the CCR, use the certification form on the State Water Board's website at http://www.swrcb.ca.gov/drinking water/certlic/drinkingwater/CCR.shtml)

Water System Name: Go		me: Golden Va	lley Municipal Water	District		
Wate	r System Nu	ımber: 1900767	57			
Furth comp	er, the syste	(date) to cum certifies that the	stomers (and approprinformation contained	iate notices of availate in the report is correct	eport was distributed on bility have been given). t and consistent with the ontrol Board, Division of	
Certified by: Name: Signature:		Name:	Frank Springer			
		Signature:	no	e-21		
		Title:	District Engineer			
		Phone Number:	(661) 248-8501	Date:	May 19, 2020	
items		and fill-in where app		methods. Specify other	r direct delivery methods	
	Good faith" efforts were used to reach non-bill paying consumers. Those efforts included t following methods:					
	Posting the CCR on the Internet at www					
	Mailing the CCR to postal patrons within the service area (attach zip codes used)					
	 Advertising the availability of the CCR in news media (attach copy of press release) Publication of the CCR in a local newspaper of general circulation (attach a copy of the published notice, including name of newspaper and date published) Posted the CCR in public places (attach a list of locations) Delivery of multiple copies of CCR to single-billed addresses serving several persons, such as apartments, businesses, and schools 					
		ivery to community er (attach a list of ot		a list of organizations)		
	For systems serving at least 100,000 persons: Posted CCR on a publicly-accessible internet site at the following address: www					
	For investo	or-owned utilities: I	Delivered the CCR to t	he California Public U	tilities Commission	
This	s form is provided	d as a convenience for use t	o meet the certification require	ement of the California Code of	f Regulations, section 64483(c).	